

# Contributions to the Psychodinae (Diptera: Psychodidae) Fauna of Turkey

*Hasan Koç*<sup>1\*</sup>, *Rudiger Wagner*<sup>2</sup>, *Alper Tonguç*<sup>1</sup>

<sup>1</sup> Muğla University, Science and Art Faculty, Department of Biology, 48170 Kötekli, Muğla, Turkey; e-mail: khasan@mu.edu.tr; alpertonguc@mu.edu.tr

<sup>2</sup> Limnologische Fluss-Station Schlitz der MPG, P.O. Box 260, D-36105 Schlitz, Deutschland; e-mail: rwagner@mpil-ploen.mpg.de.

**Abstract:** Nineteen (Psychodidae) Psychodinae species were collected between April 2003 and April 2005 in the south-west Anatolian region of Turkey. The following species were recorded for the first time in Turkey: *Satchelliella nubila* (MEIGEN, 1818), *Satchelliella pilularia* (TONNOIR, 1940), *Satchelliella trivialis* (EATON, 1893), *Pericoma bosnica* KREK, 1967, *Pericoma bunae* KREK, 1979, *Pericoma exquisita* EATON, 1893, *Pericoma motasi* VAILLANT, 1978, *Tonnoiriella pulchra* (EATON, 1893), *Psychoda minuta* (BANKS, 1894), *Psychoda parthenogenetica* TONNOIR, 1940, *Panimerus notabilis* (EATON, 1893), *Mormia malickyi* VAILLANT, 1974, *Clogmia albipunctata* (WILLISTON, 1893) and *Peripsychoda fusca* (MACQUART, 1826). The locations of 14 new collection sites from Turkey were recorded.

**Key words:** Diptera, Psychodidae, Turkey, Southwest Anatolia, new records.