

# Age Structure and Body Size Variation in Populations of *Darevskia bithynica* (Méhely, 1909) (Reptilia: Lacertidae) from Different Altitudes in North-western Turkey

Serkan Gül<sup>1\*</sup>, Nurhayat Özdemir<sup>1</sup>, Yusuf Kumlutaş<sup>2</sup>, Salih Hakan Durmuş<sup>3</sup>, Çetin Ilgaz<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Arts and Sciences, Recep Tayyip Erdoğan University, Rize, Turkey

<sup>2</sup>Department of Biology, Faculty of Science, Dokuz Eylül University, Buca-İzmir, Turkey

<sup>3</sup>Department of Biology, Faculty of Education, Dokuz Eylül University, Buca-İzmir, Turkey

**Abstract:** We present the first data of body size and age structure (based on the skeletochronological examination) of *Darevskia bithynica* from two localities characterised with different altitudes, i.e. Zonguldak (50 m) and Bolu (1200 m). Males from both lowland and highland populations were slightly larger than females. This difference was not significant for the lowland population. The maximum lifespan was found to be six years for both populations. Intersexual differences in snout-vent length were male-biased and this may be linked to the climatic conditions.

**Key words:** Age structure, *Darevskia bithynica*, sexual size dimorphism, skeletochronology, Turkey

## Introduction

The lacertid genus *Darevskia* Arribas, 1997 includes 31 species. It has a wide distribution: from Armenia, Azerbaijan, Georgia, Iran and Turkey in Western Asia, to Bulgaria, Greece, Romania, Russia, Serbia and Turkmenistan in Eastern, South-eastern and Central Europe (ARNOLD *et al.* 2007, GRECHKO *et al.* 2007, AHMADZADEH *et al.* 2013). Thirteen species of the genus *Darevskia* occur in Turkey: *D. armeniaca* (Méhely, 1909), *D. bendimahiensis* (Schmidtler, Eiselt and Darevsky, 1994), *D. bithynica* (Méhely, 1909), *D. clarkorum* (Darevsky and Vedmederja, 1977), *D. derjugini* (Nikolskij, 1898), *D. nairensis* (Darevsky, 1967), *D. parvula* (Lantz and Cyrén, 1913), *D. praticola* (Eversmann, 1843), *D. raddei* (Boettger, 1892), *D. rudis* (Bedriaga, 1886), *D. saphirina* (Schmidtler, Eiselt and Darevsky, 1994), *D. unisexualis* (Darevsky, 1966), *D. uzzelli* (Darevsky and Danielyan, 1977) and *D. valentini* (Boettger, 1892) (see ANANJEVA *et al.* 2006; BARAN *et al.* 2012; ARRIBAS *et al.* 2013). *Darevskia bithynica* has two sub-

species, the nominotypical *D. b. bithynica* (MÉHELY 1909) and *D. b. tristis* (Lantz and Cyrén, 1936), inhabiting north-west of Anatolia (ARRIBAS *et al.* 2013). It is a small lacertid with length approximately up to 75 mm. It lives mostly on rock exposures, uses crevices as refuges, and prefers habitats in and around herbaceous vegetation (ARRIBAS *et al.* 2013). Since *Darevskia bithynica* was raised to the species level by Arribas *et al.* (2013), This should be changed as “It has not yet been assessed for the IUCN Red List”

The determinations of age, growth rates and stage duration in amphibians and reptiles have relied largely on the skeletochronological method. It uses growth marks in bone tissue or lines of arrested growth (LAG, see CASTANET *et al.* 1993). Numerous scientists have applied this technique to lizards (e.g. ARAKELYAN 2002, ROITBERG and SMIRINA 2006a, 2006b, ROITBERG 2007; KUO *et al.* 2009, GUARINO *et al.* 2010, GUARINO 2010, KIM *et al.* 2010, ALTUNIŞIK *et al.* 2013, ARAKELYAN *et al.* 2013, GHARZI and YARI

\*Corresponding author: serkan.gul@erdogan.edu.tr

2013, GÜL *et al.* 2014, ÜZÜM *et al.* 2014, GÜL *et al.* 2015, ÜZÜM *et al.* 2015, YAKIN and TOK 2015).

In this study, we present the first information on the age structure, body size and lifespan of individuals of two breeding populations of *D. bithynica* from the north-west of Anatolia. We also examine altitudinal differences between populations of this species.

## Material and Methods

Seventy-four specimens of *D. bithynica* (37♂♂ and 37♀♀), deposited in the Fauna and Flora Research and Application Centre of Dokuz Eylül University (Buca-İzmir), were examined. They were from two populations (Bolu and Zonguldak) in the north-west of Anatolia, with 54 of these specimens being used only for skeletochronological analysis. Specimens were captured in mid April 2009 at altitudes of 1200 m a.s.l. (Bolu, 40°33'13.21"N, 31°15'52.68"E) and 50 m a.s.l. (Zonguldak, 41°5'58.18"N, 31°15'17.61"E).

The climate of Zonguldak varies from the coastal areas to the inland areas due to the mountains that run parallel to the coast. Summers are warm and humid, and the average temperature is around 11°C in April. Winters are cool and damp. Precipitation is the heaviest in autumn and early winter, and minimal in spring, the average precipitation is around 60.9 mm in April (MGM 2015). Bolu has cold and snowy winters, and warm summers with cool nights. The average temperature is around 9°C in April, and the average precipitation is around 51 mm in April (MGM 2015).

In order to identify sex, we used the presence/absence of hemipenes that are retracted in hemipenial sacks at the base of the tail. The snout-vent length (SVL) of all specimens was measured using a digital calliper with an accuracy of 0.01 mm. Snout-vent length was used to study sexual size dimorphism (SSD) between sexes and age groups, and to present relationship between SVL and age of *D. bithynica* specimens. LOVICH and GIBBONS (1992) proposed that SSD was quantified with the sexual dimorphism index (SDI). According to LOVICH and GIBBONS (1992),  $SDI = (\text{mean size of larger sex} / \text{mean size of smaller sex}) \pm 1$ . In this formula  $\pm 1$  is arbitrarily expressed as +1 if males are larger than females and defined as negative, or -1 if females are larger than males.

According to the standard protocol, LAGs were counted on transverse sections of the middle part of phalangeal diaphyses using a portion of the second phalanx from the third toe of the hind foot. After digits were dissected, the phalangeal bones were kept in


water for 24 h, and then were decalcified with 5% nitric acid for 2 h. Then they were kept in water for 12 h. Cross sections (18  $\mu\text{m}$ ) of the middle part of phalangeal diaphysis were prepared using a freezing microtome and stained with Ehrlich's haematoxylin. Cross sections were treated with glycerol for observation under a light microscope.

Age data were not normally distributed (Shapiro-Wilk test,  $P < 0.05$ ), whereas SVL data were normally distributed (Shapiro-Wilk test,  $P > 0.05$ ). Therefore, Mann-Whitney U test and Independent sample t-test were used to compare relationship between sexes with respect to age and SVL, correspondingly. Pearson's correlation coefficient was used to infer the pattern of relationships between age and SVL. All statistic analyses were performed using SPSS version 22 (IBM SPSS Statistics for Windows, Armonk, NY).

## Results

Cross sections of the phalangeal bones of adults showed that LAGs were pronounced enough to be counted easily (Fig. 1). Endosteal resorption appeared in almost all cross sections of adult specimens. Cross sections in which the oldest lines were difficult to distinguish because of the endosteal resorption were not included in the analyses.

Males from both lowland and highland populations were slightly larger than females (Table 1). Nevertheless, no significant difference was observed between the sexes in the lowland population (independent samples t-test,  $t = 0.777$ ,  $df = 16$ ,  $P = 0.449$ ,  $SDI = -0.025$ ). However, there was a significant dif-



**Fig. 1.** Cross section (18  $\mu\text{m}$ ) of the middle part of phalangeal diaphysis of a male specimen of *Darevskia bithynica* from the highland population (SVL: 67.3 mm). The LAGs are indicated by black arrows (m.c. = marrow cavity, e.b. = endosteal bone)

ference between the sexes in the highland population (independent samples t-test,  $t=3.861$ ,  $df=34$ ,  $P=0.000$ ,  $SDI:-0.082$ ). Snout-vent length ranged from 59.2 to 67.92 mm (average  $63.67\pm 0.62$ ) for males and from 50.94 to 66.2 mm (average  $58.87\pm 1.15$ ) for females of the highland population, and from 55.24 to 69.56 mm for males and from 54.62 to 66.12 mm for females of the lowland population. There were not significant differences between the populations for both sexes in terms of SVL (independent samples t-test, for females:  $t=1.444$ ,  $df=22$ ,  $P=0.163$ ; for males:  $t=-0.368$ ,  $df=28$ ,  $P=0.716$ ; Table 2).

In the lowland population, age ranged between four and six years in both males and females (mean:  $5.2\pm 0.44$ ;  $4.87\pm 0.44$ , respectively). In the highland population, it ranged between four and six years (mean:  $5.0\pm 0.21$ ) for males and three and six years (mean:  $4.68\pm 0.35$ ) for females. However, there was no significant difference in terms of average age between males and females both for the lowland population (Mann-Whitney U-test,  $U=32.5$ ,  $z=-0.722$ ,  $P=0.470$ ) and for the highland population (Mann-Whitney U-test,  $U=127.5$ ,  $z=-1.124$ ,  $P=0.261$ ). There were not significant differences between the populations in terms of age (Mann-Whitney U-test, for females:  $U=59$ ,  $z=-0.325$ ,  $P=0.745$ ; for males:  $U=85$ ,  $z=-0.703$ ,  $P=0.482$ ) for both sexes (Table 2). Both SVL and age were combined to determine sexual maturity of *D. bithynica* specimens. Thus, individuals of both populations were determined to have reached sexual maturity at the age of two. A correlation was not found between age and SVL in both males ( $P=0.883$ ,  $r=0.035$ ) and females ( $P=0.371$ ,  $r=0.24$ ) for the two populations.

## Discussion

Altitudinal gradients offer a natural experiment by providing divergent conditions at relatively close geographical proximity. At the highland, activity seasons are relatively short, a phenomenon especially relevant to ectotherms such as lizards (PERRY 2007). In fact, animals in the highland population and northern latitudes live longer than animals in the lowland population and southern latitudes. Moreover, animals in cooler climates from highlands and northern latitudes have larger bodies (SEARS, ANGILLETTA 2004). In addition, lizards from the lowland populations are expected to have a smaller mean body size than specimens in the highland population (ROITBERG, SMIRINA 2006b). However, our results show that female specimens from the lowland population were larger than females from the highland population, but this was not statistically significant. Male specimens

**Table 1.** Descriptive statistics of age, SVL, and SSD indexes of *Darevskia bithynica* populations

Population	SVL (mm)		Age (Years)	
	Males	Females	Males	Females
Zonguldak				
n	10	8	10	8
Min-Max	55.24-69.56	54.62-66.12	4-6	4-6
Mean	63.20	61.66	5.2	4.87
SE	1.34	1.45	0.44	0.44
SSD	-0.025			
Bolu				
n	20	16	20	16
Min-Max	59.2-67.92	50.94-66.2	4-6	3-6
Mean	63.67	58.87	5	4.68
SE	0.62	1.15	0.21	0.35
SSD	-0.082			

**Table 2.** Statistical differences between females and males in terms of SVL and age based on parametric and nonparametric tests

Sexes	SVL			Age		
	t	df	P	u	z	P
Female	1.444	22	0.163	59	-0.325	0.745
Male	-0.368	28	0.716	85	-0.703	0.482

from both populations had equal sizes (Table 1). We observed endosteal resorption in all cross sections. According to some studies (CAETANO, CASTANET 1993; ESTEBAN *et al.* 1999), this could be related to changes in the altitudinal gradient because populations living in the highland exhibit less resorption than lowland populations.

Our study points out the life history traits of *D. bithynica* across two altitudes from two different populations in the north-west of Anatolia. Our results show that the males of *D. bithynica* are larger than the females from the two populations (male-biased SSD). These results are concordant with the studies of *Lacerta agilis exigua* and *L. a. boemica* (ROITBERG 2007). Recently, it has been found that the mean SVL of *Acanthodactylus boskianus* males that inhabit South-eastern Anatolia is significantly greater than that of females (ÜZÜM *et al.*, 2014). Similarly, ALTUNIŞIK *et al.* (2013) have found that intersexual differences in body size of *Eremias strauchi strauchi* were male-biased; they have shown that males live longer than females. In addition, GÜL *et al.* (2014) have reported that males of *Darevskia rudis* from lowlands were larger than females but the difference in the highlands is not similar to our results. SSD may arise from differences in behaviour, demography, life history, physiology, ecology, age at sexual

maturity, patterns of energy use, and evolution of males and females within a population (Cox et al., 2003). Besides, intersexual variation in SSD correlates with population differences in latitude, altitude, climate, and geophysical features (COX, 2006; COX et al., 2007). For instance, Sears and ANGILLETTA (2004) have reported that cold adapted *Sceloporus graciosus* show larger SVL in individuals from populations at low altitude that undergo longer periods activity than specimens in populations at high altitude that undergo shorter periods of annual activity; they have also stated that *S. graciosus* at high altitude exhibits faster rates of growth in a relatively cool environment than lizards in lower altitudes and warmer environmental. Thus, environmental conditions in lowlands and highlands can have a significant effect on life history of *D. bithynica*. Similarly, *D. rudis* individuals from low altitude display distinctly larger mean SVL and age than from high altitudes (GÜL et al., 2014). Life history traits of *D. bithynica* across altitudinal gradi-

ents may be affected by climatic conditions because females from the highland population in Bolu (annual mean temperature 10.6°C; average amount of precipitation 45.7 mm) living in a colder and more arid environment would tend to be younger than females from the lowland population in Zonguldak (annual mean temperature 13.7°C; average amount of precipitation 101.7 mm). However, this altitudinal gradient is not statistically significant.

In conclusion, our results based on skeletochronological analysis indicated that the mean lifespan of specimens from the lowland population in both sexes is longer and intersexual differences in SVL are male-biased. However, neither differences in mean life span nor differences in mean SVL for different altitudes are statistically significant.

**Acknowledgement:** This work forms part of a project (Project No. 2009.KB.FEN.003) supported by the Scientific Research Project Coordination Unit, Dokuz Eylül University, Turkey.

## References

- ALTUNIŞIK A., Ç. GÜL, N. ÖZDEMİR, M. TOSUNOĞLU, T. ERGÜL 2013. Age Structure and Body Size of the Strauch's Racerunner, *Eremias strauchi strauchi* KESSLER 1878. – *Turkish Journal of Zoology*, **37**: 539-543.
- AHMADZADEH F., M. FLECKS, M. A. CARRETERO, O. MOZAFFARI, W. BÖHME, D. J. S. HARRIS, FREITAS, D. RÖDDER 2013. Cryptic speciation patterns in Iranian Rock Lizards uncovered by integrative taxonomy. – *Plos One*, **8**: 1-17.
- ANANIEVA N. B., N. L. ORLOV, R. G. KHALIKOV, I. S. DAREVSKY, I. S. RYABOV, A. V. BARABANOV 2006. An Atlas of the Reptiles of North Eurasia. Taxonomic, Diversity, Distribution, Conservation Status. Sofia, Bulgaria (Pensoft Series Faunistica 47), 247 p.
- ARAKELYAN M. 2002. The study of age, growth, and longevity in the triploid hybrids of rock lizards of the genus *Darevskia* in Armenia. – *Russian Journal of Herpetology*, **9**: 63-68.
- ARAKELYAN M., R. PETROSYAN, Ç. ILGAZ, Y. KUMLUTAŞ, S. H. DURMUŞ, Y. TAYHAN, F. DANIELYAN 2013. A skeletochronological study of parthenogenetic lizards of genus *Darevskia* from Turkey. – *Acta Herpetologica*, **8**: 99-104.
- ARNOLD E. N., O. ARRIBAS, S. CARRANZA 2007. Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. – *Zootaxa*, **1430**: 1-86.
- ARRIBAS O. J., Ç. ILGAZ, Y. KUMLUTAŞ, S. H. DURMUŞ, A. AVCI, N. ÜZÜM 2013. External morphology and osteology of *Darevskia rudis* (BEDRIAGA 1886), with a taxonomic revision of the Pontic and Small-Caucasus populations (Squamata: Lacertidae). – *Zootaxa*, **3626**: 401-428.
- BARAN İ., Ç. ILGAZ, A. AVCI, Y. KUMLUTAŞ, K. OLGUN 2012. Türkiye Amfibi ve Sürüngenleri. Ankara, Turkey (TÜBİTAK Popüler Bilim Kitapları), 204 p. (In Turkish).
- CAETANO M. H., J. CASTANET 1993. Variability and micro evolutionary patterns in *Triturus marmoratus* from Portugal: Age, size longevity, and individual growth. – *Amphibia-Reptilia*, **14**: 117-129.
- CASTANET J., H. FRANCILLON-VIEILLOT, F. J. MEUNIER, A. DE RICQLE 1993. Bone and individual aging. – In: HALL, B. K. (Ed.): Bone Growth B, Vol: 7, Boca Raton: Florida, 245-283.
- COX R. M., S. L. SKELLY, H. B. JOHN-ALDER 2003. A comparative test of adaptive hypotheses for sexual size dimorphism in lizards. – *Evolution*, **57**: 1653-1669.
- COX R. M. 2006. A test of the reproductive cost hypothesis for sexual size dimorphism in Yarrow's spiny lizard, *Sceloporus jarrovi*. – *Journal of Animal Ecology*, **75**: 1361-1369.
- COX R. M., M. A. BUTLER, H. B. JOHN-ALDER 2007. The evolution of sexual size dimorphism in reptiles. – In: Fairbairn D. J., W. U. Blanckenhorn, T. Székely (Eds.): Sex, Size and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism. Oxford, United Kingdom (Oxford University Press), 38-49.
- GHAZZI A., A. YARI 2013. Age determination in the Snake-eyed Lizard, *Ophisops elegans*, by means of skeletochronology (Reptilia: Lacertidae). – *Zoology in the Middle East*, **59**: 10-15.
- GRECHKO V. V., A. A. BANNIKOVA, S. A. KOSUSHKIN, N. L. RYABININA, K. D. MILTO, I. S. DAREVSKY, D. A. KRAMEROV 2007. Molecular genetic diversification of the lizard complex *Darevskia raddei* (Sauria: Lacertidae): Early stages of speciation. – *Molecular Biology*, **41**: 764-775.
- GUARINO F. M. 2010. Structure of the femora and autotomous (postpygal) caudal vertebrae in the three-toed skink *Chalcides chalcides* (Reptilia: Squamata: Scincidae) and its applicability for age and growth rate determination. – *Zoologischer Anzeiger*, **248**: 273-283.
- GUARINO F. M., I. D. GIÀ, R. SINDACO 2010. Age and growth of the sand lizards (*Lacerta agilis*) from a high Alpine population of northwestern Italy. – *Acta Herpetologica*, **5**: 23-29.
- GÜL S., N. ÖZDEMİR, Y. KUMLUTAŞ, Ç. ILGAZ 2014. Age structure and body size in three populations of *Darevskia rudis* (BEDRIAGA 1886) from different altitudes. – *Herpetozoa*, **26**: 151-158.

- GÜL S., N. ÖZDEMİR, A. AVCI, Y. KUMLUTAŞ, Ç. ILGAZ 2015. Altitudinal effects on life history of Anatolian lizard (*Apathya cappadocica*, WERNER 1902) from the southeastern Anatolia, Turkey. – *Turkish Journal of Zoology*, **39**: 507-512.
- KIM J. K., J. Y. SONG, J. H. LEE, D. PARK 2010. Physical characteristics and age structure of Mongolian racerunner (*Eremias argus*; Lacertidae). – *Journal of Ecology and Field Biology*, **33**: 325-331.
- KUO C. Y., Y. T. LIN, Y. S. LIN 2009. Sexual size and shape dimorphism in an Agamid lizard, *Japalura swinhonis* (Squamata: Lacertilia: Agamidae). – *Zoological Studies*, **48**: 351-61.
- LOVICH J. E., J. W. GIBBONB 1992. A review of techniques for quantifying sexual size dimorphism. – *Growth, Development, and Aging*, **56**: 269-281.
- MGM (General Directorate of Meteorology), 2015. İllerimize ait istatistiki veriler (Yearly Data). <<http://www.mgm.gov.tr/veridegerlendirme/il-ve-ilceler-istatistik.aspx?m=ZONGULDAK>> (accessed 14.05.15) (InTurkish).
- PERRY G. 2007. Movement patterns in lizards: measurement, modality, and behavioral correlates. – In: REILLY S., L. McBRAYER, D. B. MILES (Eds.): *Lizard Ecology: The Evolutionary Consequences of Foraging Mode*. Cambridge, United Kingdom (Cambridge University Press), 13-48.
- ROITBERG E. S., E. M. SMIRINA 2006a. Age, body size and growth of *Lacerta agilis boemica* and *L. strigata* (Reptilia, Lacertidae): a comparative study of two closely related lizard species based on skeletochronology. – *Herpetological Journal*, **16**: 133-148.
- ROITBERG E. S., E. M. SMIRINA 2006b. Adult body length and sexual size dimorphism in *Lacerta agilis boemica* (Reptilia, Lacertidae): between-year and inter locality variation. – In: CORTI C., P. LO CASCIO, M. BIAGGINI (Eds.): *Mainland and Insular Lizards: A Mediterranean Perspective*. Florence, Italy (Florence University Press), 175-187.
- ROITBERG E. 2007. Geographic variation in sexual size dimorphism within a widespread lizard species. – In: FAIRBAIRN D. J., W. U. BLANCKENHORN, T. SZEKELY (Eds.): *Sex, Size and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism*. Oxford, United Kingdom (Oxford University Press), 143-153.
- SEARS M. W., M. J. ANGILLETTA 2004. Body size clines in Sceloporus lizards: Proximate mechanisms and demographic constraints. – *Integrative and Comparative Biology*, **44**: 433-442.
- ÜZÜM N., Ç. ILGAZ, Y. KUMLUTAŞ, Ç. GÜMÜŞ, A. AVCI 2014. The body size, age structure, and growth of Bosc's fringe-toed lizard, *Acanthodactylus boskianus* (DAUDIN 1802). – *Turkish Journal of Zoology*, **38**: 383-388.
- ÜZÜM N., A. AVCI, Y. KUMLUTAŞ, N. BEŞER, Ç. ILGAZ 2015. The first record of age structure and body size of the Suphan Racerunner, *Eremias suphani* BAŞOĞLU, HELLMICH 1968. – *Turkish Journal of Zoology*, DOI: 10.3906/zoo-1408-39.
- YAKIN B. Y., C. V. TOK 2015. Age Estimation of *Anatololacerta anatolica* (WERNER 1902) in the Vicinity of Çanakkale, by Skel-etchronology. – *Turkish Journal of Zoology*, **39**: 66-73.

Received: 13.02.2015

Accepted: 29.09.2015

