



The Dipterans (Insecta: Diptera) of the Bulgarian Black Sea Coast

Zdravko Hubenov¹

¹National Museum of Natural history, Bulgarian Academy of Sciences, 1 Tsar Osvoboditel Blvd., 1000 Sofia, Bulgaria;
E-mail: zkhubenov@abv.bg; zhubenov@nmnhs.com

Academic Editor: *Dragan Chobanov*

Abstract: Up until now, a total of 1556 dipteran species of 82 families have been reported from the Bulgarian Black Sea Coast. The families Cecidomyiidae, Chironomidae, Syrphidae, Chloropidae and Tachinidae (101–162 species) are the most numerous. The families of Ephydriidae (70.4 % of all Bulgarian species presented), Chloropidae (67.5 %), Tabanidae (55.3 %), Calliphoridae (46.3%) and Cecidomyiidae (44.4 %) show the biggest relative species richness in the Black Sea Coast area. Recorded low relative species richness is often due to lack of sufficient investigations, such as for the families Tipulidae, Mycetophilidae and Phoridae. The faunistic similarity between the different parts of the coast is scrutinised. The dipterans belong to 116 areographical categories, divided into 2 groups: 1) species with a Mediterranean type of distribution (191 species – 12.3%) – more thermophilic and distributed mainly in the southern parts of the Palaearctic; 2) species with Palaearctic and Eurosiberian type of distribution (1365 species – 87.7%) – more eurybiontic and widely distributed in the Palaearctic. The distribution of the zoogeographical categories in the different parts of the coast is presented. A total of 30 (1.9%) endemics have been established. Of the reported species, 40 are plant pests and 63 have a medical importance.

Key words: Diptera, Bulgarian Black Sea Coast, faunistic composition, distribution, zoogeography

Introduction

During the last 70 years, the Black Sea Coast has been under significant anthropogenic impact and landscape changes. The dynamic nature of the fauna and its conservation require periodic updating of the faunistic diversity of the taxonomic groups. The present study covers, for the first time, all dipteran families from the Black Sea Coast of Bulgaria.

The first data on Diptera from the Bulgarian Black Sea Coast were reported by Löw (1862). Later, the works of Nedelkov (1912), Szilády (1934), Zilahi (1934), Jacentkovský (1936, 1937), Valkanov (1936)

and Drensky (1939, 1942) appeared, which contained data from the coast. After World War II, a large number of articles related to dipterans of the Black Sea Coast were published. Studies on Chironomidae (Cvetkov 1955a, 1955b, 1957, 1958, Michailova 1973a, 1973b, 1974, 1975, 1980, 1982, 1989), Culicidae (Božkov 1961, 1966, 1991, Božkov et al. 1969, Christova et al. 1971, Mikov 2005, 2008, Mikov et al. 2013), Cecidomyiidae and Sciaridae (Skuhrava et al. 1991, 1992, Dimitrova & Mohrig 1993) predominate and separate families of Brachycera from the coastal habitats (Beschovski 1964a, 1964b, 1965, 1966a, 1966b, 1972a, 1972b, 1973a, 1973b, 1975a,

1975b, 1976, 1978, 1993, 1994, 1996, Lavčiev 1969, 1980, 2003, Zielke 1916a, 1916b, 2018, 2019). Some applied entomological studies that are not related to the coast include faunistic data. The hydrobiological studies of the coastal water bodies have also had an important faunistic contribution (Valkanov 1948, 1954, 1957, Caspers 1951a, 1951b, 1952, Stoichev 1996, 1998, Vassilev & Konsulov 1998, Kovachev et al. 1999, Georgiev & Nikolov 2010, Varadinova et al. 2010, Todorova 2014, Hubenov et al. 2015). Volumes from the Fauna of Bulgaria series and catalogues have been published (Beschovski 1985, 2009, 2013, Lavčiev 2003, Bechev 2006, 2010, Kechev et al. 2020, Hubenov 2021a, Kechev 2021), which include data from the Bulgarian Black Sea Coast. The studies of some foreign authors also contain information about the coast of Bulgaria (Slípka 1959, Hradský & Moucha 1964, 1967, Bankowska 1967a, 1967b, Starý 1982, 2009, Krzemiński 1984, Mendl 1986, Wagner & Joost 1988, Krzemiński & Starý 1989, Povolný & Verves 1990, Starý & Krzemiński 1993, Ježek et al. 2020, Černý et al. 2022, Lutovinovas et al. 2022, Dvořák et al. 2024).

The aim of this work is to present the fauna, distribution and zoogeography of dipterans from the Bulgarian Black Sea Coast.

Materials and methods

The Bulgarian Black Sea Coast (Fig. 1) stretches for 380 km from the border with Romania at Cape Sivriburun to the Rezovska River at the border with Turkey. The width of this territory varies from 5 to 60 km. It is divided into 5 parts (Galabov 1966, Daneva & Mishev 1979, Krastev 2002): **Dobrudzha** (from the border with Romania to the Batova River), **Varna** (between the valleys of the Batova and Kamchia Rivers), **Stara Planina** (between the valleys of the Kamchia and Hadzhiyska Rivers), **Burgas** (from the foot of the Eminska Planina Mts. and the Hadzhiyska River to Medni Rid) and **Strandzha** (south of the Burgas Lowland to the Rezovska River) districts. The Black Sea Coast includes lowlands below 100 m a.s.l., plateaus and plains up to about 350 m a.s.l. and hilly lands up to about 600 m a.s.l. In the Black Sea coastal zone, the altitudinal belt up to 300 m a.s.l. prevails. The Bulgarian Black Sea Coast belongs to the Continental-Mediterranean climatic zone, with the Mediterranean climatic influence being stronger in the Burgas and Strandzha regions (Sabev & Stanev 1959, Tichkov 1976, Stanev 1991, Velev 2002, 2010). The Mediterranean-Black Sea precipitation regime is presented. There are about 40 lakes, marshes and river spills along the coast (Varbanov




Fig. 1. Partition of the Black Sea Coast: 1) Dobrudzha district, 2) Varna district, 3) Stara Planina district, 4) Burgas district, 5) Strandzha district

2002, Hristova 2012). The lakes are of the firth and lagoon type. Nineteen rivers enter the Black Sea, and their mouth areas have a firth character. The Black Sea region is included in the Euxinian Province of the European deciduous forest area and in the zone of the xerothermic oak forests (Stojanov 1966, Velchev et al. 1982, Bondev 1991, 2002, Velchev 2002). The Eastern Stara Planina and Strandzha Mts. are rich in Euxinian elements. The vegetation along the coast contains Mediterranean elements with typical communities around rivers being the flooded (longose) forests. Specific psammophyte vegetation is developed on the sand dunes. Characteristic halophyte communities are represented on the saline soils. The coastal strip with a width of 5 to 40 km forms the Black Sea zoogeographical region (Georgiev 1978, 2002, Hubenov 2021b). There is a Mediterranean influence on the fauna, which increases to the south. The Mediterranean elements in some halophilic and halobiontic insects reach 72%. Caspian relicts are concentrated in the coastal firth lakes and river mouths. A specific fauna is connected with the hyperhaline lakes.

The material had been collected after 1890 from 261 localities. The main part of it is stored in

the National Museum of Natural History and the Institute of Biodiversity and Ecosystem Research. All dipteran species reported from the Bulgarian Black Sea Coast are included. For a number of widespread species the authors did not give the localities and mentioned they occur everywhere. In some publications, the localities are presented generalised for the entire coast. A number of foreign publications contain data on the Bulgarian Black Sea Coast.

Weaknesses of the literature data that limit the obtaining of an equivalent information when comparing taxonomic groups include: different levels of study of the separate families; lack of exact localities for some of the reported species; insufficient data for the vertical distribution; existence of rich synonymy; outdated data; lack of generalised investigations; unexplored parts of the coast; prolonged periods of data accumulation; priority studies on groups of economic and medical importance. Some old data for Chironomidae collected during the hydrobiological studies are problematic since determination of the larvae is not always possible.

The classification of the dipterans is based on the works of Soós & Pap (1984-1993), McAlpine (1989), Wood & Borkend (1989), Woodley (1989), Nagatomi (1996), Papp & Darvas (1997, 1998, 2000a, 2000b), Yeates & Wiegmann (1999), Nartshuk (2003), Ziegler (2003), Oosterbroek (2006), Yeates et al. (2007) and Pape et al. (2011).

Areographical analysis for the taxa categorization was used. This method allows obtaining data information about species complexes with different zoogeographical character based on the published data regarding species distribution and results of the faunistic research. These complexes contain zoogeographical information about the taxonomic groups which, combined with the origin of the ranges, determines the zoogeographical character of the fauna. The distribution of the species according to the zoogeographical categories in the different coastal parts is scrutinised. The classification of the areas is based on the works of Geptner (1936), Darlington (1957), Kryzhanovskiy (1965, 1976, 2002), de Lattin (1967), Emeljanov (1974, 2018), Müller (1974, 1980), Udvardi (1975), Crosskey & White (1977), Malicky et al. (1983), Gorodkov (1984), Grehan (1988, 1993), Lopatin (1989), Vigna Taglianti et al. (1999), Procheş & Ramdhani (2012), Holt et al. (2013) and Ficetola et al. (2017). The border between the Western and Eastern Palaearctic along the Yenisei River is accepted. To compare the fauna, Czekanowski-Dice-Sørensen coefficient of similarity is used (Czekanowski 1909, Dice 1945, Sørensen 1948).

Results

A total of 1556 dipteran species of 82 families have been recorded from the Bulgarian Black Sea Coast so far (Table 1), which accounts for 30.0% of the 5190 species, known from Bulgaria – Nematocera 395 species (25.4%) and Brachycera 1161 species (74.6%). The families Tachinidae (162 species – 10.4%), Cecidomyiidae (116 species – 7.5%), Chloropidae (112 species – 7.2%), Syrphidae (110 species – 7.1%) and Chironomidae (101 species – 6.5%) are the most numerous. The remaining families contain less than 100 species. Of the largest families (with over 200 species in Bulgaria), Limoniidae, Mycetophilidae, Cecidomyiidae, Chironomidae, Dolichopodidae, Phoridae, Syrphidae, Agromysidae, Muscidae and Tachinidae, between 0.9% and 44.4% of the Bulgarian species have been reported from the Black Sea Coast. The families Mycetophilidae and Phoridae, widely represented in other parts of the country, have not been studied along the coast. It is expected that a significant number of species of the Tachinidae family will additionally be established for the Black Sea Coast. The families Culicidae, Stratiomyidae, Tabanidae, Chamaemyiidae, Sciomyzidae, Chloropidae and Ephydriidae are best represented – between 55.3% and 74.5% of the Bulgarian species. The families Culicidae, Ceratopogonidae, Chironomidae, Chloropidae, Ephydriidae and Muscidae have been studied longest and the most systematically in the whole country. Along the Black Sea Coast, they comprise between 30.8% and 74.5% of the Bulgarian species. Some smaller families are well represented, of which between 30% and 75.0% of the taxa found in Bulgaria have been established along the coast. Of the infraorders, Orthorrhapha includes 250 species (16.1% of the coastal species and 28.3% of the Bulgarian species), related to the better study of Stratiomyidae (60.0% of the Bulgarian species), Tabanidae (55.3%), Asilidae (28.7%) and Dolichopodidae (30.7%). Significantly more species include Cyclorrhapha (911 species – 58.5% of the coastal species and 36.6% of the Bulgarian species). This infraorder combines 52 families of the Black Sea Coast, some of which (Tethinidae, Canacidae and Ephydriidae) are connected with the coastal habitats and others are rich in species. The diverse habitats of the coast support a rich dipteran fauna, which is insufficiently studied. The number of the established species is probably about 50-60% of the real species composition of the Bulgarian Black Sea Coast.

The wide distribution of Diptera suggests that thorough research would reveal similar species in different regions. Most dipterans have vast ranges

Table 1. Distribution of the dipterans (Insecta: Diptera) of the Bulgarian Black Sea Coast. The number in brackets after the families give the number of the established species in Bulgaria. In the second column, the figure after the brackets indicates the percentage from the Bulgarian species.

Families	Total number	Zoning of the Bulgarian Black Sea Coast				
		Dobroudzha	Varna	Stara Planina	Bourgas	Strandzha
NEMATOCERA (1674)	395 (25.4) 23.6	94 (19.0)	279 (32.5)	46 (19.7)	129 (21.4)	110 (19.1)
Tipulomorpha (343)	45 (2.9) 13.1	3 (0.6)	19 (2.2)	12 (5.1)	4 (0.7)	31 (5.4)
Tipulidae (89)	4 (0.3) 4.5	3 (0.6)	4 (0.5)	3 (1.3)	3 (0.5)	3 (0.5)
Limoniidae (221)	41 (2.6) 18.6		15 (1.7)	9 (3.9)	1 (0.2)	28 (4.8)
Bibionomorpha (687)	178 (11.4) 25.9	27 (5.4)	161 (18.7)	16 (6.9)	14 (2.3)	27 (4.7)
Bibionidae (13)	2 (0.1) 15.4	1 (0.2)	2 (0.2)	1 (0.4)	1 (0.2)	1 (0.2)
Mycetophilidae (262)	14 (0.9) 5.3	3 (0.6)	9 (1.0)			4 (0.7)
Ditomyiidae (3)	1 (0.06) 33.3		1 (0.1)			
Bolitophilidae (10)	1 (0.06) 10.0					1 (0.2)
Diadocidiidae (3)	1 (0.06) 33.3		1 (0.1)			
Keroplataidae (48)	15 (1.0) 31.2		9 (1.0)	2 (0.9)		8 (1.4)
Sciaridae (84)	28 (1.8) 33.3	1 (0.2)	26 (3.0)			
Cecidomyiidae (261)	116 (7.5) 44.4	22 (4.4)	113 (13.2)	13 (5.6)	13 (2.2)	13 (2.3)
Psychodomorpha (109)	7 (0.4) 6.4	1 (0.2)	2 (0.2)	1 (0.4)	5 (0.8)	
Psychodidae (102)	6 (0.4) 5.9	1 (0.2)	2 (0.2)	1 (0.4)	4 (0.7)	
Trichoceridae (7)	1 (0.06) 20.0				1 (0.2)	
Ptychopteromorpha (5)	1 (0.06) 20.0		1 (0.1)			
Ptychopteridae (6)	1 (0.06) 16.6		1 (0.1)			
Culicomorpha (521)	164 (10.5) 31.5	63 (12.7)	96 (11.2)	17 (7.3)	106 (17.6)	52 (9.0)
Chaoboridae (1)	1 (0.06) 100.0	1 (0.2)				
Culicidae (47)	35 (2.2) 74.5	19 (3.8)	25 (2.9)	12 (5.1)	29 (4.8)	25 (4.3)
Ceratopogonidae (66)	27 (1.7) 40.9		16 (1.9)		25 (4.2)	1 (0.2)
Chironomidae (328)	101 (6.5) 30.8	43 (8.7)	55 (6.4)	5 (2.1)	52 (8.6)	26 (4.5)
BRACHYCERA (3506)	1161 (74.6) 33.1	402 (81.0)	580 (67.5)	187 (80.3)	473 (78.6)	467 (80.9)
ORTHORRHAPHA (881)	250 (16.1) 28.3	65 (13.1)	154 (17.9)	16 (6.9)	97 (16.1)	55 (9.5)
Stratiomyomorpha (51)	30 (1.9) 58.8	3 (0.6)	23 (2.7)	2 (0.9)	11 (1.8)	11 (1.9)
Stratiomyidae (50)	30 (1.9) 60.0	3 (0.6)	23 (2.7)	2 (0.9)	11 (1.8)	11 (1.9)
Tabanomorpha (107)	51 (3.3) 47.7	20 (4.0)	31 (3.6)	7 (3.0)	27 (4.5)	15 (2.6)
Rhagionidae (18)	3 (0.2) 16.7		1 (0.1)			3 (0.5)
Athericidae (2)	1 (0.06) 50.0			1 (0.4)		
Tabanidae (85)	47 (3.0) 55.3	20 (4.0)	30 (3.5)	6 (2.6)	27 (4.5)	12 (2.1)
Asilomorpha (221)	68 (4.4) 30.8	13 (2.6)	43 (5.0)	4 (1.7)	29 (4.8)	14 (2.4)
Acroceridae (3)	1 (0.06) 33.3				1 (0.2)	
Bombyliidae (89)	28 (1.8) 31.5	8 (1.6)	24 (2.8)	3 (1.3)	13 (2.2)	2 (0.3)
Mythicomyiidae (2)	1 (0.06) 50.0				1 (0.2)	1 (0.2)
Therevidae (10)	5 (0.3) 50.0	1 (0.2)	4 (0.5)			
Asilidae (115)	33 (2.1) 28.7	4 (0.8)	15 (1.7)	1 (0.4)	14 (2.3)	11 (1.9)
Empidomorpha (502)	101 (6.5) 20.1	29 (5.8)	57 (6.6)	3 (1.3)	30 (5.0)	15 (2.6)
Empididae (183)	14 (0.9) 7.6	3 (0.6)	6 (0.7)		3 (0.5)	4 (0.7)
Hybotidae (87)	19 (1.2) 21.8	5 (1.0)	8 (0.9)		7 (1.2)	3 (0.5)
Microphoridae (5)	1 (0.06) 20.0					1 (0.2)
Dolichopodidae (225)	67 (4.3) 30.7	21 (4.2)	43 (5.0)	3 (1.3)	20 (3.3)	7 (1.2)
CYCLORRHAPHA (2491)	911 (58.5) 36.6	337 (67.9)	426 (49.6)	171 (73.4)	379 (63.0)	412 (71.4)
ASCHIZA (565)	119 (7.6) 21.1	20 (4.0)	42 (4.9)	16 (6.9)	50 (8.3)	70 (12.1)
Lonchopteridae (4)	2 (0.1) 50.0	1 (0.2)	2 (0.2)	1 (0.4)	1 (0.2)	1 (0.2)
Phoridae (230)	2 (0.1) 0.9	1 (0.2)				1 (0.2)
Pipunculidae (26)	5 (0.3) 19.2		4 (0.5)		1 (0.2)	1 (0.2)
Syrphidae (304)	110 (7.1) 36.2	18 (3.6)	36 (4.2)	15 (6.4)	48 (8.0)	67 (11.6)
SCHIZOPHORA (1925)	792 (50.9) 41.1	317 (63.9)	384 (44.7)	155 (66.5)	329 (54.7)	342 (59.3)
Acalyptrata (941)	431 (27.7) 45.8	199 (40.1)	261 (30.4)	40 (17.2)	208 (34.6)	142 (24.6)
Psilidae (7)	1 (0.06) 14.3		1 (0.1)			

Table 1. Continuation.

Families	Total number	Zoning of the Bulgarian Black Sea Coast				
		Dobroudzha	Varna	Stara Planina	Bourgas	Strandzha
Conopidae (41)	16 (1.0) 39.0	1 (0.2)	10 (1.2)		3 (0.5)	5 (0.9)
Lonchaeidae (5)	1 (0.06) 20.0				1 (0.2)	
Otitidae (13)	8 (0.5) 61.5	1 (0.2)	6 (0.7)		1 (0.2)	
Platystomatidae (4)	1 (0.06) 25.0		1 (0.1)			
Tephritidae (79)	32 (2.1) 40.5	7 (1.4)	29 (3.4)	2 (0.9)	10 (1.7)	4 (0.7)
Piophilidae (3)	1 (0.06) 33.3	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Lauxaniidae (23)	5 (0.3) 13.0		2 (0.2)			6 (1.0)
Chamaemyiidae (29)	19 (1.2) 65.5	14 (2.8)	11 (1.3)		8 (1.3)	8 (1.4)
Coelopidae (3)	2 (0.1) 66.7	1 (0.2)	2 (0.2)		1 (0.2)	
Dryomyzidae (2)	1 (0.06) 50.0				1 (0.2)	
Sciomyzidae (37)	23 (1.5) 62.2	3 (0.6)	16 (1.9)		4 (0.7)	5 (0.9)
Helcomyzidae (1)	1 (0.06) 100.0	1 (0.2)	1 (0.1)			
Sepsidae (11)	1 (0.06) 9.0					1 (0.2)
Odiinidae (2)	1 (0.06) 50.0					1 (0.2)
Agromyzidae (309)	36 (2.3) 11.6	3 (0.6)	4 (0.5)	6 (2.6)	19 (3.2)	14 (2.4)
Opomyzidae (7)	5 (0.3) 71.4		4 (0.5)		2 (0.3)	2 (0.3)
Anthomyzidae (5)	1 (0.06) 20.0		1 (0.1)			
Perisclididae (1)	1 (0.06) 100.0					1 (0.2)
Asteiidae (3)	2 (0.1) 66.7	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Carnidae (8)	3 (0.2) 37.7	1 (0.2)	2 (0.2)	1 (0.4)	2 (0.3)	1 (0.2)
Tethinidae (7)	7 (0.4) 100.0	6 (1.2)	3 (0.3)	3 (1.3)	5 (0.8)	4 (0.7)
Canacidae (1)	1 (0.06) 100.0	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Milichiidae (8)	5 (0.3) 62.5	2 (0.4)	2 (0.2)		5 (0.8)	1 (0.2)
Chloropidae (166)	112 (7.2) 67.5	68 (13.7)	71 (8.3)	9 (3.9)	68 (11.3)	38 (6.6)
Siphonellopsidae (2)	2 (0.1) 100.0		2 (0.2)			
Heleomyzidae (51)	11 (0.7) 21.6	2 (0.4)	6 (0.7)	1 (0.4)	3 (0.5)	2 (0.3)
Trioxselididae (1)	1 (0.06) 100.0	1 (0.2)				
Sphaeroceridae (60)	20 (1.3) 33.3	12 (2.4)	12 (1.4)	2 (0.9)	6 (1.0)	6 (1.0)
Curtonotidae (1)	1 (0.06) 100.0		1 (0.1)		1 (0.2)	
Camillidae (1)	1 (0.06) 100.0	1 (0.2)			1 (0.2)	
Drosophilidae (17)	11 (0.7) 64.7	2 (0.4)	2 (0.2)		3 (0.5)	8 (1.4)
Campichoetidae (2)	1 (0.06) 50.0	1 (0.2)	1 (0.1)			
Diastatidae (4)	2 (0.1) 50.0	2 (0.4)	2 (0.2)			
Ephydriidae (135)	95 (6.1) 70.4	67 (13.5)	66 (7.7)	13 (5.6)	61 (10.1)	32 (5.5)
Calyptrata (985)	361 (23.2) 36.6	118 (23.8)	123 (14.3)	115 (49.4)	121 (20.1)	200 (34.7)
Hippoboscidae (13)	4 (0.2) 30.8	2 (0.4)	2 (0.2)	2 (0.9)	2 (0.3)	4 (0.7)
Nycteribiidae (7)	7 (0.4) 100.0					7 (1.2)
Scathophagidae (11)	1 (0.06) 9.1		1 (0.1)		1 (0.2)	
Anthomyiidae (53)	14 (0.9) 26.4	9 (1.8)	8 (0.9)	8 (3.4)	10 (1.7)	12 (2.1)
Fanniidae (27)	9 (0.6) 33.3	3 (0.6)	4 (0.5)	3 (1.3)	3 (0.5)	8 (1.4)
Muscidae (267)	98 (6.3) 36.7	20 (4.0)	48 (5.6)	9 (3.9)	40 (6.6)	55 (9.5)
Calliphoridae (41)	20 (1.3) 46.3	7 (1.4)	9 (1.0)	5 (2.1)	18 (3.0)	9 (1.6)
Sarcophagidae (122)	39 (2.5) 32.0	17 (3.4)	17 (2.0)	7 (3.0)	20 (3.3)	12 (2.1)
Rhinophoridae (8)	1 (0.06) 12.5		1 (0.1)			
Oestridae (2)	2 (0.1) 100.0	1 (0.2)	2 (0.2)	1 (0.4)	2 (0.3)	1 (0.2)
Hypodermatidae (4)	1 (0.06) 25.0	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Gasterophilidae (4)	3 (0.2) 75.0	3 (0.6)	3 (0.3)	3 (1.3)	3 (0.5)	3 (0.5)
Tachinidae (436)	162 (10.4) 37.2	55 (11.1)	27 (3.1)	76 (32.6)	21 (3.5)	88 (15.2)
SPECIES	1556	496 (31.9)	859 (55.2)	233 (15.0)	602 (38.7)	577 (37.1)
FAMILIES	82	53 (64.6)	66 (80.5)	37 (45.1)	56 (68.3)	56 (68.3)

Table 2. Similarity of the dipteran fauna in the different parts of the Bulgarian Black Sea Coast. Common species are given in brackets

Zoning	Varna	Stara Planina	Bourgas	Strandzha
Dobroudzha	45.0 (305)	38.8 (131)	50.3 (276)	40.8 (215)
Varna		27.2 (141)	45.7 (334)	35.3 (250)
Stara Planina			35.1 (137)	36.4 (134)
Bourgas				43.0 (249)

and endemics are poorly represented [30 species – 1.9% (Nematocera – 0.4% and Brachycera – 1.5%)]. Often these are newly described taxa or rare species with unclear ranges. The distribution of Diptera along the Bulgarian Black Sea Coast is related to the specific natural conditions of the respective regions, the peculiarities of the families and the level of their study. This is evident when comparing the established species between the separate regions of the coast. They have not been evenly studied in terms of families, as separate families are well studied in some regions and almost unexplored in others. Thus, Sciaridae is represented only in the Varna district, Ceratopogonidae – around Varna and Burgas, Dolichopodidae is almost absent in the Stara Planina district, Sciomyzidae prevails in the Varna district, most Agromyzidae are found in the Burgas district and Tachinidae – from the Strandzha district. Culicidae, Chironomidae (excluding the Stara Planina district) and Ephydriidae are most evenly represented in the separate parts of the coast. The families studied more extensively throughout the years by various research groups are evenly represented along the coast. Investigations on the synovil and synanthropic forms of Tabanidae, Fanniidae, Muscidae, Calliphoridae and Sarcophagidae have enriched the faunistic composition of the Brachycera in the Black Sea Coast. Many families have not been studied and the taxa reported are from accidentally collected materials. These families are represented by widely distributed species, which do not allow for the drawing of conclusions about their presence in the separate parts of the coast. Single taxa and fragmentary data have been reported. Most material has been collected around the popular scientific and resort centres. With respect to most families, the regions of Varna and Burgas are better studied (over 40% of the species found), while the Stara Planina Mts. part is poorly studied (15.0% of the species known). When considering Brachycera and Nematocera in general (Table 1), the differences between the separate regions reach 13.5%.

Of the established dipterans, 63 species have a medical significance, and 40 species are pests on the vegetation. Of the suborder Nematocera, 22 species

have a human or veterinary medical significance, and 20 species are pests on forestry or agriculture. Most representatives with medical significance include the family Culicidae (12), followed by Ceratopogonidae (6) and Psychodidae (4). The majority of vegetation pests belong to the family Cecidomyiidae (16). Only 4 species are shared out among Tipulidae (2), Bibionidae (1) and Sciaridae (1). Of the suborder Brachycera, 41 species have a human or veterinary medical significance, and 20 species are pests on forestry or agriculture. Most representatives of medical significance include Muscidae (10) and Tabanidae (8), followed by Sarcophagidae (5), Calliphoridae (4), Gastrophilidae (3), Fanniidae (3), Hippoboscidae (3), Oestridae (2), Piophilidae (1), Anthomyiidae (1) and Hypodermatidae (1). Most vegetation pests belong to Anthomyiidae (5), Tephritidae (4) and Chloropidae (4). The families Psilidae, Lonchaeidae, Agromyzidae, Opomyzidae, Heleomyzidae, Drosophilidae and Ephydriidae are presented with one species each.

The zoogeographical categorization of the species is made on the basis of current data about their distribution. Thus, the dipterans are divided into 116 areographical categories, combined in 3 main groups and 6 complexes (Table 3).

Species distributed in the Palaearctic and beyond it. This group (470 species – 30.2%) includes 38 categories, of which 27 combine species of northern type (widely distributed in the Holarctic and Palaearctic) and eleven – species of southern type (distributed in the southern parts of the Palaearctic). In the Super Palaearctic complex, the Holarctic species are best represented [137 species – 8.8% (from 6.9% in the Stara Planina district to 9.9% in the Dobrudzha district)]. Of the other areographical categories, the Palaearctic-Oriental species [73 species – 4.7% (10 – 31 species, from 3.6% to 5.0% in the separate districts)] and the Holarctic-Oriental [41 species – 2.6% (9 – 27 species, from 2.2% to 4.7%)] forms are better represented. Among dipterans, there is a significant number of synanthropic and synbovillous forms with a cosmopolitan or subcosmopolitan distribution. Their ranges are structured and affected by the development of

Table 3. Zoogeographical characteristic of Diptera of the Bulgarian Black Sea Coast

Classification of the areas	Total number	Zoning of the Bulgarian Black Sea Coast				
		Dobrudzha	Varna	Stara Planina	Burgas	Strandzha
Species distributed in Palaearctic and out of it	470 (30.2)	198 (39.9)	234 (27.2)	95 (40.8)	215 (35.7)	222 (38.5)
NORTH TYPE	439 (28.2)	186 (37.5)	225 (26.2)	89 (38.2)	197 (32.7)	208 (36.0)
Cosmopolitan	29 (1.9)	21 (4.2)	21 (2.4)	18 (7.7)	20 (3.3)	21 (3.6)
Semicosmopolitan	17 (1.1)	11 (2.2)	11 (1.3)	9 (3.9)	11 (1.8)	10 (1.7)
Holarctic-Paleotropical-Neotropical	3 (0.2)	1 (0.2)	2 (0.2)		2 (0.3)	1 (0.2)
Holarctic-Paleotropical-Australian	5 (0.3)	4 (0.8)	3 (0.3)	3 (1.3)	3 (0.5)	5 (0.9)
Holarctic-Paleotropical	4 (0.3)	2 (0.4)	1 (0.1)		1 (0.2)	1 (0.2)
Holarctic-Neotropical-Afrotropical	2 (0.1)	2 (0.4)				1 (0.2)
Holarctic-Neotropical-Oriental	13 (0.8)	8 (1.6)	11 (1.3)	5 (2.1)	7 (1.2)	8 (1.4)
Holarctic-Afrotropical-Australian	2 (0.1)	2 (0.4)			1 (0.2)	
Holarctic-Oriental-Australian	5 (0.3)		2 (0.2)		2 (0.3)	1 (0.2)
Holarctic-Afrotropical	6 (0.4)	4 (0.8)	4 (0.5)	2 (0.9)	4 (0.7)	4 (0.7)
Holarctic-Oriental	41 (2.6)	12 (2.4)	19 (2.2)	9 (3.9)	21 (3.5)	27 (4.7)
Holarctic-Neotropical	10 (0.6)	5 (1.0)	5 (0.6)	3 (1.3)	6 (1.0)	6 (1.0)
Holarctic-Australian	8 (0.5)	2 (0.4)	5 (0.6)	1 (0.4)	3 (0.5)	3 (0.5)
Holarctic	137 (8.8)	49 (9.9)	69 (8.0)	16 (6.9)	56 (9.3)	56 (9.7)
Palaearctic-Paleotropical-Neotropical	1 (0.06)				1 (0.2)	1 (0.2)
Palaearctic-Paleotropical-Australian	8 (0.5)	4 (0.8)	5 (0.6)	2 (0.9)	5 (0.8)	6 (1.0)
Palaearctic-Afrotropical-Neotropical	1 (0.06)		1 (0.1)			
Palaearctic-Afrotropical-Australian	1 (0.06)	1 (0.2)			1 (0.2)	1 (0.2)
Palaearctic-Oriental-Australian	3 (0.2)	2 (0.4)	1 (0.1)		1 (0.2)	1 (0.2)
Palaearctic-Paleotropical	25 (1.6)	13 (2.6)	15 (1.7)	4 (1.7)	12 (2.0)	8 (1.4)
Palaearctic-Afrotropical	15 (1.0)	8 (1.6)	9 (1.0)	3 (1.3)	7 (1.2)	6 (1.0)
Palaearctic-Oriental	73 (4.7)	24 (4.8)	31 (3.6)	10 (4.3)	26 (4.3)	29 (5.0)
Palaearctic-Australian	1 (0.06)	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
West Palaearctic-Afrotropical	13 (0.8)	6 (1.2)	5 (0.6)		6 (1.0)	5 (0.9)
West Palaearctic-Oriental	8 (0.5)	1 (0.2)	3 (0.3)			4 (0.7)
West Palaearctic-Neotropical	1 (0.06)	1 (0.2)				
Disjunct Palaearctic-Oriental	7 (0.4)	2 (0.4)	1 (0.1)	3 (1.3)		2 (0.3)
SOUTH TYPE	31 (2.0)	12 (2.4)	9 (1.0)	6 (2.6)	18 (3.0)	14 (2.4)
South Palaearctic-Paleotropical-Australian	2 (0.1)	2 (0.4)	1 (0.1)	2 (0.9)	2 (0.3)	1 (0.2)
Southwest Palaearctic-Neotropical-Afrotropical-Australian	1 (0.06)					1 (0.2)
South Palaearctic-Paleotropical	8 (0.5)	5 (1.0)	3 (0.3)	1 (0.4)	4 (0.7)	2 (0.3)
South Palaearctic-Afrotropical	1 (0.06)				1 (0.2)	
Southwest Palaearctic-Afrotropical	3 (0.2)		1 (0.1)		1 (0.2)	1 (0.2)
South Palaearctic-Oriental	5 (0.3)	2 (0.4)	1 (0.1)		1 (0.2)	2 (0.3)
Southwest Palaearctic-Oriental	3 (0.2)	1 (0.2)	1 (0.1)	1 (0.4)	3 (0.5)	2 (0.3)
Afrotropical-Mediterranean	5 (0.3)	1 (0.2)	1 (0.1)	1 (0.4)	4 (0.7)	3 (0.5)
Oriental-Mediterranean	1 (0.06)					1 (0.2)
Oriental-East Mediterranean	1 (0.06)				1 (0.2)	
European-Neotropical-Oriental	1 (0.06)	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Species with Palaearctic distribution	1086 (69.8)	298 (60.1)	625 (72.8)	126 (54.1)	387 (64.3)	351 (60.8)
PALAEARCTIC TYPE	489 (31.4)	170 (34.3)	273 (31.8)	77 (33.0)	204 (40.6)	195 (33.8)
Holopalaearctic	17 (1.1)	11 (2.2)	13 (1.5)	6 (2.6)	15 (2.5)	10 (1.7)
Transpalaearctic	109 (7.0)	41 (8.3)	56 (6.5)	16 (6.9)	49 (8.1)	55 (9.5)
West and Central Palaearctic	68 (4.4)	26 (5.2)	48 (5.6)	14 (6.0)	28 (4.7)	21 (3.6)
West Palaearctic	92 (5.9)	36 (7.3)	48 (5.6)	16 (6.9)	37 (6.1)	39 (6.8)
Disjunct Palaearctic	19 (1.2)	3 (0.6)	11 (1.3)	4 (1.7)	5 (0.8)	5 (0.9)
South Palaearctic	2 (0.1)		2 (0.2)		1 (0.2)	1 (0.2)
Southwest Palaearctic	6 (0.4)	1 (0.2)	3 (0.3)	1 (0.4)	1 (0.2)	2 (0.3)
Eurosiberian-Anatolian-Central Asian	4 (0.3)		2 (0.2)		1 (0.2)	1 (0.2)
Eurosiberian-Central Asian	28 (1.8)	5 (1.0)	13 (1.5)	5 (2.1)	10 (1.5)	10 (1.7)

Table 3. Continuation.

Classification of the areas	Total number	Zoning of the Bulgarian Black Sea Coast				
		Dobrudzha	Varna	Stara Planina	Burgas	Strandzha
West Eurosiberian-Central Asian	4 (0.3)	3 (0.6)	2 (0.2)	1 (0.4)	2 (0.3)	
West Eurosiberian-West Central Asian	1 (0.06)		1 (0.1)			
West Eurosiberian-Iran-Turanian	2 (0.1)		1 (0.1)		2 (0.3)	1 (0.2)
West Eurosiberian-Anatolian-Iranian	1 (0.06)					1 (0.2)
West Eurosiberian-Anatolian-Turanian	1 (0.06)		1 (0.1)			
West Eurosiberian-Turanian	1 (0.06)	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
West Eurosiberian-Anatolian	3 (0.2)	3 (0.6)	1 (0.1)		1 (0.2)	1 (0.2)
European-Anatolian-North African	18 (1.2)	2 (0.4)	6 (0.7)	1 (0.4)	13 (2.2)	7 (1.2)
European-North African	53 (3.4)	19 (3.8)	36 (4.2)	5 (2.1)	22 (3.7)	20 (3.5)
European-Anatolian-Central Asian	7 (0.4)	3 (0.6)	4 (0.5)	1 (0.4)		2 (0.3)
European-Central Asian	7 (0.4)	3 (0.6)	2 (0.2)		2 (0.3)	2 (0.3)
European-West Central Asian	9 (0.6)	2 (0.4)	4 (0.5)	1 (0.4)	4 (0.7)	2 (0.3)
European-Southwest Asian	13 (0.8)	3 (0.6)	4 (0.5)	2 (0.9)	3 (0.5)	5 (0.9)
European-Anatolian-Iran-Turanian	5 (0.3)	3 (0.6)	3 (0.3)	2 (0.9)	4 (0.7)	3 (0.5)
European-Iran-Turanian	1 (0.06)	1 (0.2)			1 (0.2)	1 (0.2)
European-Anatolian-Iranian	7 (0.4)	2 (0.4)	4 (0.5)		2 (0.3)	2 (0.3)
European-Anatolian-Turanian	1 (0.06)					1 (0.2)
European-Iranian	2 (0.1)		1 (0.1)			1 (0.2)
European-Turanian	8 (0.5)	2 (0.4)	6 (0.7)	1 (0.4)	1 (0.2)	1 (0.2)
Species distributed in one subregion	597 (38.4)	128 (25.8)	352 (41.0)	49 (21.0)	183 (30.4)	156 (27.0)
EUROSIBERIAN TYPE	407 (26.2)	82 (16.5)	250 (29.1)	31 (13.3)	93 (15.4)	94 (16.3)
Holoeurosiberian	23 (1.5)	9 (1.8)	10 (1.2)	3 (1.3)	12 (2.0)	6 (1.0)
Transeurosiberian	13 (0.8)	3 (0.6)	7 (0.8)		2 (0.3)	2 (0.3)
West and Central Eurosiberian	26 (1.7)	9 (1.8)	16 (1.9)		11 (1.8)	8 (1.4)
West Eurosiberian	26 (1.7)	5 (1.0)	13 (1.5)	2 (0.9)	7 (1.2)	7 (1.2)
Disjunct Eurosiberian	34 (2.2)	4 (0.8)	15 (1.7)	6 (2.6)	6 (1.0)	9 (1.6)
European and South Siberian	5 (0.3)	1 (0.2)	2 (0.2)			3 (0.5)
European-Anatolian	19 (1.2)	2 (0.4)	15 (1.7)	1 (0.4)	3 (0.5)	5 (0.9)
European	217 (13.9)	42 (8.5)	144 (16.8)	14 (6.0)	42 (7.0)	47 (8.1)
East European-Siberian	1 (0.06)		1 (0.1)			
East European	3 (0.2)	1 (0.2)	2 (0.2)			1 (0.2)
Central and East European	2 (0.1)	1 (0.2)	1 (0.1)			
Central and South European-Anatolian	7 (0.4)		2 (0.2)	1 (0.4)	3 (0.5)	4 (0.7)
Central and South European	20 (1.3)	3 (0.6)	14 (1.6)	4 (1.7)	3 (0.5)	2 (0.3)
Central and Southeast European	11 (0.7)	2 (0.4)	8 (0.9)		4 (0.7)	
MEDITERRANEAN TYPE	160 (10.3)	41 (8.3)	83 (9.7)	15 (6.4)	83 (13.8)	55 (9.5)
Mediterranean and South Siberian	1 (0.06)					1 (0.2)
North Mediterranean and South Far East	1 (0.06)	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Mediterranean-Central Asian	5 (0.3)	1 (0.2)			4 (0.7)	4 (0.7)
Mediterranean-West Central Asian	8 (0.5)	2 (0.4)	5 (0.6)	2 (0.9)	5 (0.8)	4 (0.7)
Mediterranean-Iran-Turanian	6 (0.4)	1 (0.2)	2 (0.2)		3 (0.5)	1 (0.2)
Mediterranean-Iranian	7 (0.4)	2 (0.4)	5 (0.6)		5 (0.8)	3 (0.5)
Mediterranean-Turanian	2 (0.1)		1 (0.1)	1 (0.4)	1 (0.2)	
East Mediterranean-Central Asian	1 (0.06)				1 (0.2)	1 (0.2)
East Mediterranean-Iran-Turanian	1 (0.06)				1 (0.2)	
North Mediterranean-West Central Asian	2 (0.1)	1 (0.2)	1 (0.1)		1 (0.2)	2 (0.3)
North Mediterranean-Iranian	3 (0.2)		2 (0.2)		2 (0.3)	1 (0.2)
Northeast Mediterranean-Iranian	1 (0.06)					1 (0.2)
Southeast European-West Central Asian	1 (0.06)	1 (0.2)				1 (0.2)
Central and South European-Iran-Turanian	2 (0.1)	2 (0.4)	2 (0.2)	1 (0.4)	2 (0.3)	2 (0.3)
Central and Southeast European-Iranian	3 (0.2)	1 (0.2)		1 (0.4)		1 (0.2)
Central (Middle) and South European-Turanian	3 (0.2)	2 (0.4)	2 (0.2)		1 (0.2)	1 (0.2)

Table 3. Continuation.

Classification of the areas	Total number	Zoning of the Bulgarian Black Sea Coast				
		Dobrudzha	Varna	Stara Planina	Burgas	Strandzha
Central and Southeast European-Lebanonian	1 (0.06)	1 (0.2)				
Central and South European-Anatolian-North African	1 (0.06)				1 (0.2)	
Central and South European-North African	5 (0.3)	2 (0.4)	1 (0.1)		2 (0.3)	1 (0.2)
South European-North African	20 (1.3)	8 (1.6)	14 (1.6)	1 (0.4)	13 (2.2)	3 (0.5)
Southeast European-Iranian	1 (0.06)		1 (0.1)			1 (0.2)
South European-Turanian	1 (0.06)					1 (0.2)
Holomediterranean	24 (1.5)	3 (0.6)	11 (1.3)	3 (1.3)	12 (2.0)	8 (1.4)
East Mediterranean	7 (0.4)	1 (0.2)	5 (0.6)	2 (0.9)	2 (0.3)	1 (0.2)
North Mediterranean	9 (0.6)	3 (0.6)	6 (0.7)		3 (0.5)	3 (0.5)
South European	15 (1.0)	2 (0.4)	8 (0.9)		9 (1.5)	3 (0.5)
Northeast Mediterranean	1 (0.06)	1 (0.2)			1 (0.2)	
Southeast European-Anatolian	7 (0.4)	2 (0.4)	5 (0.6)		3 (0.5)	4 (0.7)
Southeast European	16 (1.0)	3 (0.6)	7 (0.8)	1 (0.4)	9 (1.5)	4 (0.7)
Balkan-Caucasian-Iranian	1 (0.06)		1 (0.1)	1 (0.4)		
Balkan-Caucasian	1 (0.06)	1 (0.2)	1 (0.1)	1 (0.4)	1 (0.2)	1 (0.2)
Balkan-Anatolian	3 (0.2)		2 (0.2)			1 (0.2)
ENDEMIC	30 (1.9)	5 (1.0)	19 (2.2)	3 (1.3)	7 (1.2)	7 (1.2)
Balkan subendemic	1 (0.06)		1 (0.1)			
Balkan endemic	10 (0.6)	3 (0.6)	6 (0.7)	3 (1.3)	4 (0.7)	2 (0.3)
Bulgarian endemic	15 (1.0)	2 (0.4)	11 (1.3)		2 (0.3)	2 (0.3)
Regional endemic	4 (0.3)		1 (0.1)		1 (0.2)	3 (0.5)
SPECIES	1556	496	859	233	602	577

the human civilization (before the contemporary research). These species most often have a high abundance, present in diverse habitats and increase the percentage of the Super Palaearctic taxa in the non-specialized studies.

Species distributed within the Palaearctic region. The group (1086 species – 69.8%) includes from 126 to 625 species (from 54.1% to 72.8%) in the separate parts of the Bulgarian Black Sea Coast. It is divided into species distributed only in the Palaearctic, but in more than one subregion and species distributed within one Palaearctic subregion.

Species distributed only in the Palaearctic but in more than one subregion (Palaearctic type). Taxa, whose areas include more than one Palaearctic subregion in latitudinal direction, belong to this group. In total, 28 areographical categories are represented in the dipteran fauna of the Bulgarian Black Sea Coast (Table 3), which combine 489 species (31.4%). The character of the Palaearctic complex is determined by the Transpalaearctic [109 species – 7.0% (from 6.5% to 9.5% in the separate coastal areas)], West Palaearctic [92 species – 5.9% (from 5.6% to 7.3%)], West and Central Palaearctic [68 species – 4.4% (from 3.6% to 6.0%)] and European-North African [53 species – 3.4% (from 2.1%

to 4.2%)] species. The Eurosiberian-Central Asian (28 species), European-Anatolian-North African (18 species), Holopalaearctic and Disjunct Palaearctic (17 – 19 species) taxa are well represented. The correlation of the mentioned categories is consistent in the separate vegetation belts and varies from 0.4% to 9.5% (1 to 56 species). Seventeen species (1.2%), from 3 to 11 species in the separate coastal regions have longitudinal disjunction of the areas with regard to Siberia and Central Asia. Probably, some of these species are presented with sparse populations and will be studied in more details as a result of further research. Most often, a latitudinal disjunction of the areas of this complex is lacking (Gorodkov 1984, Josifov 1988, Hubenov 2015).

Species distributed within one subregion of the Palaearctic. This group (597 species – 38.4%) includes from 49 to 352 species (from 21.0% to 41.0%) in the separate parts of the Bulgarian Black Sea Coast (Table 3). The group combines species with Eurosiberian and Mediterranean types of distribution (50 categories), as well as endemics. The Mediterranean-Central Asian species are also included here, following Kryzhanovskiy (1965, 2002) and Lopatin (1989), who combine the Mediterranean and Central Asian subregions.

The Eurosiberian species include 14 areo-geographical categories (407 species or 26.2%). These are from 31 to 250 species (from 13.3% to 29.1%) in the separate parts of the Bulgarian Black Sea Coast. The European [217 species (13.9%) – from 14 to 144 species (from 6.0% to 16.8%) in the separate parts of the coast], Disjunct Eurosiberian [34 species (2.2%) – from 4 to 15 species (from 0.8% to 2.6%)], West and Central Eurosiberian [26 species (1.7%) – from 0 to 16 species (from 0% to 1.9%)] and West Eurosiberian [26 species (1.7%) – from 2 to 13 species (from 0.9% to 1.5%)] taxa are the most numerous. The Holoeurosiberian, European-Anatolian, Central and South European, Transeurosiberian and Central and Southeast European species are well represented. The Eurosiberian complex includes 34 species with disjunctive areas – a longitudinal disjunction for Siberia and Central Asia and latitudinal disjunction with boreomontane or boreoalpine distribution (Gorodkov 1984; Josifov 1988; Hubenov 2015). In other systematic groups (such as Cerambycidae of Coleoptera), the boreomontane and montane species connected with the coniferous communities descend below 1000 m a.s.l. due to the large afforestations of conifers in the lowlands (Georgiev & Hubenov 2006).

The Mediterranean species include 32 areo-geographical categories (160 species or 10.3%). These are from 1 to 24 species (0.06% to 1.5%) in the separate parts of the Bulgarian Black Sea Coast (Table 3). They are better represented in the southern parts of the coast (Burgas and Strandzha districts). Because of the big variety of their areas, the group is divided into subgroups with different origin, distribution and ecological peculiarities of the taxa. This complexity contributes to establishing of various zoogeographical classifications of Bulgaria (Josifov 1981, 1986, 1988, 1999, Gruev 1988, 1995, 2000a, 2000b, 2000c, 2002, Heiss & Josifov 1990, Gruev & Kusmanov 1994, Hubenov 1996, 2008, Gruev & Bechev 2000, Popov 2002). For the separate areo-geographical categories, the difference between the regions of the Bulgarian Black Sea Coast is the largest in the South European-North African species (1.8%). The Holo-mediterranean (24 species – 1.5%), South European-North African (20 species – 1.3%), Southeast European (16 species – 1.0%) and South European (15 species – 1.0%) taxa are the most numerous.

Endemics. This category includes taxa, which are not distributed outside the Balkan Peninsula. The percentage of endemism is low in Diptera (30 species – 1.9%). The endemism is differently presented in Nematocera (7 species – 0.4%) and Brachycera (23 species – 1.5%). The Bulgarian (15 species – 1.0%) and Balkan (10 species – 0.6%) endemic

forms prevail. The greatest number of endemics (19 species – 2.2%) have been found in the Varna district of the coast. The Burgas district is significantly poorer in endemic taxa (7 species – 1.2%). Local endemics have not been established among Diptera of the Bulgarian Black Sea Coast. Some of the endemic dipterans (9 species) were described after 1990 or are rare taxa with unclear range.

Discussion

The faunistic similarity among the different parts of the Bulgarian Black Sea Coast (Table 2) is related to their specific natural conditions and insufficient research. A higher similarity (from 45.0% to 50.2%) is noticed among the low parts of the coast – Burgas, Varna and Dobrudzha districts. The similarity between these territories is greater than their similarity with the neighboring Stara Planina and Strandzha districts (from 27.2% to 43.0%). The poor research of the Stara Planina coast and the small number of common species probably determine the low similarity between the Varna, Stara Planina and Burgas parts of the coast (from 27.2% to 35.1%). The better research of the Varna and Burgas districts and the many common species increase their faunistic similarity (45.7%), although they are not neighboring territories. Further investigations of the separate parts of the Bulgarian Black Sea coast would likely confirm them to be similar in terms of the species composition of Diptera. This is related both to the natural habitats and the wide distribution of the dipterans, their high mobility and low level of endemism.

The difference (%) between the separate districts for Super Palaearctic species reaches 13.6% and varies from 27.2% (Varna district) to 40.8% (Stara Planina district). For the separate areo-geographical categories, this difference is the highest for the the cosmopolitan (5.3%) and subcosmopolitan (2.6%) species. The establishment of other species of the group in the district of the Stara Planina coast is very likely, owing to their distribution and insufficient studies of this part of the Bulgarian Black Sea Coast. It is accepted that the taxa of northern type have the vast ranges and ecological flexibility. The species of southern type (31 species – 2.0%) are better represented in the southern parts of the coast (Burgas and Strandzha districts). They are not important for the zoogeographical characteristic due to their small number (6 – 18 species, from 1.0% to 3.0%). Usually, the Super Palaearctic group is scantily presented and is not determinant for the zoogeographical characteristic of the most taxa in the

Bulgarian terrestrial fauna (with the exception of the coastal fauna). In the highly mobile forms (such as Diptera), the group is well represented and exceeds 25% (Hubenov 2015, 2019, 2025a, 2025b). Along the Bulgarian Black Sea Coast, it is represented (%) better in the Stara Planina and Dobrudzha parts of the coast (Table 3).

Species from the Palearctic complex are well represented in the high mobile groups and comprise about 20-30% of the species. The difference between the separate coastal districts (from 77 to 273 species) reaches 8.8% and varies from 31.8% (Varna district) to 40.6% (Burgas district) of the species. For the separate areographical categories, this difference is largest for the Transpalearctic species (3.0%). The vast areas and wide vertical distribution of the taxa of this complex are an indication of the greater ecological flexibility of its species. The dominance of the Palearctic complex is related to the insufficient studies of these territories and the non-systematic sampling, resulting in the collection of more common and widespread species (Hubenov 2019).

The ratio of the areographical categories is different for the separate families: in some cases, the Holoeurosiberian, Disjunct Eurosiberian and European species are almost equal in number, with a total of 50% Eurosiberian forms, while in other families the Central and South European species are better represented or in general, the Eurosiberian taxa are poorly represented. The number of species by areographical categories varies in the separate coastal regions from 0% to 16.8% (0-144 species). For the separate categories, this difference is greatest in the European taxa (10.8%). The greatest number of Eurosiberian species has been established in the Varna district of the coast (250 species – 29.1%). In all other regions of the Bulgarian Black Sea Coast, taxa with Super Palearctic and Palearctic ranges dominate. Probably, some “Eurosiberian” species actually could be Palearctic taxa. The Bulgarian Black Sea Coast has specific climatic conditions, which are connected with the presence of the Eurosiberian species. These climatic conditions are well expressed on the eastern slopes of the Stara Planina and Strandzha Mts., where an inversion of the vegetation is observed - the oak forests are above beech forests (Bondev 1991, 2002, Velchev 2002).

Most Mediterranean species (22 species – 68.5%) are found in two or three regions of the coast. The significant percentage of these species and their relatively scarce populations are due to the lower ecological flexibility of the Mediterranean forms in comparison with the previous ones. There are no significant differences in the distribution of the

well presented areographical categories among the separate parts of the coast. When comparing them, a higher percentage of the Mediterranean forms southwards is noticed (from 8.3% to 13.8%), which is connected with the geographical location of the regions and the strengthening of the Mediterranean influence south of the Stara Planina Mts. This does not apply to the Stara Planina Mts. coast (6.4%), due to its poor research (only 15 Mediterranean species have been established – 6.4% of the all Mediterranean forms). The percentage of the Mediterranean species is slightly higher than the average one for Bulgaria (9.1% – Hubenov 2021a) and significantly higher than the percentage for the mountains (from 3.3% to 9.2% – Hubenov 2017, 2019, 2025).

Conclusion

This study reveals that 1556 dipteran species belonging to 82 families are so far recorded along the Bulgarian Black Sea Coast, representing 30.0% of the 5190 dipteran species known from Bulgaria. The most species-rich families are Tachinidae (162 species; 10.4%), Cecidomyiidae (116 species; 7.5%), Chloropidae (112 species; 7.2%), Syrphidae (110 species; 7.1%), and Chironomidae (101 species; 6.5%). All remaining families are represented by fewer than 100 species. The highest relative species richness in the Black Sea Coast region is observed in the families Ephydriidae (70.4% of the Bulgarian fauna), Chloropidae (67.5%), Tabanidae (55.3%), Calliphoridae (46.3%), and Cecidomyiidae (44.4%). Conversely, low relative species richness recorded for some families is likely attributable to insufficient research effort, particularly in Tipulidae, Mycetophilidae, and Phoridae.

The dipteran fauna can be divided into 2 main groups: 1) species with Mediterranean type of distribution (191 species – 12.3%) – more thermophilic and distributed mainly in the southern parts of the Palearctic and the lower parts of the mountains. The species of the southern type, distributed in the Palearctic and beyond it, are related to this group as well; 2) species with Palearctic and Eurosiberian type of distribution (1365 species – 87.7%) – more eurybiontic and widely distributed in the Palearctic. The species of the northern type, distributed in the Palearctic and beyond it, are related to this group as well. The zoogeographical character of Diptera is determined by the second group.

The percentage of species with a Mediterranean type of distribution of the Bulgarian Black Sea coast is 2.1% higher than the average for Bulgaria (10.2%).

Acknowledgements: The author would like to thank Dr. Theo Zeegers for providing data on the species of the family Tachinidae (Supplementary material – *) collected with Malaise traps by Dr. Cees van Achterberg in 1998.

Supplementary material. Table. Species composition and distribution of the dipterans (Insecta: Diptera) of the Bulgarian Black Sea Coast

Excel Spreadsheet:

<https://acta-zoologica-bulgaria.eu/2026/002996.xls>.

References

- Bankowska R. 1967a. Matériaux pour l'étude des diptères de Bulgarie (Stratiomyidae, Conopidae, Pipunculidae et Acroceridae). *Fragmenta Faunistica* (Warszawa) 13 (17): 303-314.
- Bankowska R. 1967b. Matériaux pour l'étude des Syrphidae (Diptera) de Bulgarie. *Fragmenta Faunistica* (Warszawa), 13 (21): 345-389.
- Bechev D. 2006. The fungus gnats of the families Bolitophilidae, Diadocidiidae, Ditomyiidae and Keroplatidae (Diptera: Sciaroidea) of Bulgaria. *Scientific Studies of the University of Plovdiv, Biology, Animalia* 42: 21-83.
- Bechev D. 2010. On the family Mycetophilidae (Insecta: Diptera) in Bulgaria. *ZooNotes, Supplement 1*, 72 pp.
- Beschovski V. 1964a. Diptera aus dem litoralen Gebiet der bulgarischen Schwarzmeerküste. *Bulletin de l'Institut de Pisciculture et de Pêcherie* (Varna) 4: 91-98. (In Bulgarian with German summary).
- Beschovski V. 1964b. Beitrag zum Studium der halobionten und halophilen Diptera Brachycera vom Bulgarischen Küstenbereich des Schwarzen Meeres. *Zoologischer Anzeiger* 172 (4): 261-264.
- Beschovski V. 1965. The Insects of the Bulgarian Black Sea Coast. Varna, Darzhavno izdatelstvo. 96 p. (In Bulgarian).
- Beshovski V. 1966a. Several families of Diptera unknown to Bulgarian fauna: Dryomyzidae, Chamaemyiidae, Trichoscelidae, Astiidae, Canaceidae and Milichiidae. *Bulletin de l'Institut de Zoologie et Musée* 21: 11-13. (In Bulgarian, English summary).
- Beschovski, V. 1966b. *Allotrichoma valcanovi* n. sp. – eine neue Art von (Diptera, Ephydridae) aus Bulgarien. *Comptes Rendus de l'Académie Bulgare des Sciences* 19 (9): 851-853.
- Beschovski V. 1972a. A Contribution to the dipteran Fauna (Diptera, Brachycera) in the Bulgarian Black Sea Coast. II. *Bulletin de l'Institut de Zoologie et Musée* 34: 5-14. (In Bulgarian, English summary).
- Beschovski V. 1972b. Ecologic Investigations on the representatives of Diptera-Brachycera from the salt biotops along the Bulgarian Seaside. (Sum.). I. Communication: Dipteran Fauna from the Cliffs. *Bulletin de l'Institut de Zoologie et Musée* 35: 5-21. (In Bulgarian, English summary).
- Beschovski V. 1973a. Contribution to the Dipterous Fauna (Diptera-Brachycera) of the Bulgarian Black Sea Coast. III. *Bulletin de l'Institut de Zoologie et Musée* 37: 33-41. (In Bulgarian with English summary).
- Beschovski V. 1973b. *Hilara psammophytophila* n. sp. (Diptera: Empididae) – a new species for the Bulgarian Black Sea Coast. *Comptes Rendus de l'Académie Bulgare des Sciences* 26 (8): 1061-1063.
- Beschovski V. 1975a. The Black Sea Coast inundated by waves and its Dipterous fauna (Diptera-Brachycera). *Hydrobiologia* 2: 3-18. (In Bulgarian with English summary).
- Beschovski V. 1975b. Ökologische Untersuchungen über Vertreter von Diptera-Brachycera von den Salzbiotopen der bulgarischen Schwarzmeerküste. IV. Diptera-Brachycera from the salt grounds. *Ecology* 1: 64-74. (In Bulgarian with German summary).
- Beschovski V. 1976. Diptera-Brachycera in the dunes along the Bulgarian Black Sea coast. In: *Terrestrial fauna of Bulgaria. Materials*. Sofia, Bulgarian Academy of Sciences. pp. 35-59. (In Bulgarian with English summary).
- Beschovski V. 1978. Faunistic and taxonomic investigations on the genus *Chlorops* Meigen, 1803 (Diptera, Chloropidae) from Bulgaria with description of a new subgenus. *Nouv. Rev. Entomol.* 8: 397-402. Bozhkov D. 1961. Mückenlarven aus dem Gebiet von Burgas und dem Strandzagebirge. In: Paspalev G. (Ed.). *Prirodni ognishta na zaraza v Strandzha planina*. Sofia, Bulgarian Academy of Sciences. pp. 147-153. (In Bulgarian with German summary).
- Beschovski V. 1985. Fauna bulgarica. 14. Grass flies (Diptera, Chloropidae). Sofia, Editio Academica. 220 pp. (In Bulgarian).
- Beschovsky V. 1993. The species of the genus *Lipara* Meigen (Diptera, Chloropidae), and its galls on the common reed along the Bulgarian Black Sea coast. Second National Scientific Conference of Entomology, Sofia, 25-27 October, 1993: 14-18. (In Bulgarian with English summary).
- Beschovski V. 1994. Contribution to the study of the West Palearctic Tethinidae (Diptera). *Acta Zoologica Bulgarica*, 47: 16-29.
- Beschovski V. 1996. Qualitative composition and distribution of the Ephydridae species (Diptera) on the moist zones of the Shablenska like system. *Acta Entomologica Bulgarica* 2 (1): 16-20. (Bulgarian; English summary).
- Beschovski V. 2009. Fauna bulgarica. 28. Insecta: Diptera: Ephydridae, Tethinidae, Canacidae. Sofia, Editio Academica "Professor Marin Drinov". 423 pp. (In Bulgarian).
- Beschovski V. 2013. *Catalogus faunae bulgaricae*. 8. Carnoidea. Insecta: Diptera Brachycera Acaliptratae, Sofia, Editio Academica "Professor Marin Drinov". 152 pp.
- Bondev I. 1991. The vegetation of Bulgaria. Sofia: Kliment Ohridski University Press. 184 pp. (In Bulgarian with English summary).
- Bondev I. 2002. Geobotanic regioning. In: Copralev I. (Ed.). *Geography of Bulgaria*. Sofia, ForCom, 336-352. (In Bulgarian with English summary).
- Bozhkov D. 1966a. Krovososushtie komary Bolgarii [Blood-sucking mosquitoes of Bulgaria]. *Entomologicheskoe Obozrenie* 45 (3): 570-574. (In Russian).
- Bozhkov D. 1991. Kravosmucheshtie komari [The blood-sucking mosquitoes]. Sofia, Bulgarian Academy of Sciences. 223 pp. (In Bulgarian).
- Bozhkov D., Christova T. & Canev I. 1969. Stechmücken an der bulgarischen Schwarzmeerküste. *Bulletin de l'Institut de Zoologie et Musée* 29: 151-166. (In Bulgarian with German summary).
- Caspers H. 1951a. Quantitative Untersuchungen über die Bodentierwelt des Schwarzen Meeres im bulgarischen Küstenbereich. *Archiv für Hydrobiologie*, 45 (1-2): 1-192.
- Caspers H. 1951b. Biozönotische Untersuchungen über die Stran-

- darthropoden im bulgarischen Küstenbereich des Schwarzen Meeres. *Hydrobiologia*, den Haag, 3 (2): 131-193.
- Caspers H. 1952. Untersuchungen über die Tierwelt von Meeres-salinen an der bulgarischen Küste des Schwarzen Meeres. *Zoologischer Anzeiger*, 148 (5/8): 243-259.
- Černý M., Barták M., Kubík Š. & Vála M. 2022. New records of Agromyzidae (Diptera) from Bulgaria. *Zootaxa* 5175 (4): 401-438. <https://doi.org/10.11646/zootaxa.5175.4.1>
- Christova T., Canev I. & Smilova D. 1971. Promeni vav vidoviya sastav na kravosmucheshtite komari ot bulgarskoto Chernomorsko kraybrezhie [Changes in the species composition of the blood-sucking mosquitoes from the Bulgarian Black Sea coast]. *Letopisi na HEI* 5 (31): 173-183. (In Bulgarian).
- Crosskey R. W. & White G. B. 1977. The Afrotropical Region. A recommended term in zoogeography. *Journal of Natural History*, 11: 541-544.
- Cvetkov (Zvetkov) L. 1955a. Die Chironimidenfauna der Bulgarischen Schwarzenmeereen. *Bulletin de l'Institut zoologique de l'Academie des Sciences de Bulgarie* 4/5: 215-249. (In Bulgarian with German summary).
- Cvetkov L. 1955b. Untersuchungen über das Futter der Fische im Beloslavsko-See. *Bulletin de l'Institut zoologique de l'Academie des Sciences de Bulgarie* 4/5: 329-352. (In Bulgarian with German summary).
- Cvetkov L. 1957. Zoobenthos des Beloslav-Sees. *Bulletin de l'Institut zoologique de l'Academie des Sciences de Bulgarie* 6: 381-439. (In Bulgarian with German summary).
- Cvetkov (Tsvetkov) L. 1958. Investigations into the microbenthos of the Bulgarian Black Sea lakes. *Bulletin de l'Institut zoologique de l'Academie des sciences de Bulgarie*, 7: 219-250. (In Bulgarian with English summary).
- Czekanowski J. 1909. Zur Differentiadiagnose der Neandertalgruppe. *Korrespondenzblatt der deutschen Gesellschaft für Anthropologie, Ethnologie und Urgeschichte* 40: 44-47.
- Darlington P. 1957. *Zoogeography*. New York, Wiley. 675 pp.
- Daneva M. & Mishev K. 1979. Prirodni teritorialni kompleksi (landshaftni kompleksi). [Natural territorial complexes (landscape complexes)]. In: Mishev K. & Popov V. *Bulgarian Black Sea Coast*. Sofia, Bulgarian Academy of Sciences, 89-114. (In Bulgarian).
- Dice L. 1945. Measures of the amount of ecologic association between species. *Ecology* 26 (3): 297-302.
- Dimitrova B. & Mohrig W. 1993. Beitrag zur Trauermückenfauna Bulgariens (Diptera, Sciaridae). II. *Acta Zoologica Bulgarica* 46: 89-96.
- Drensky K. 1939. The varieties of *Anopheles maculipennis* Meigen and their relation to the distribution of malaria. *Mitteilungen der Bulgarischen Entomologischen Gesellschaft in Sofia* 10: 31-44. (In Bulgarian with English summary).
- Drensky P. 1942. Über die Insektenfauna des Küstengebietes nördlich von Warna. *Mitteilungen der Bulgarischen Entomologischen Gesellschaft in Sofia* 12: 15-44. (In Bulgarian with German summary).
- Dvořák L., Dvořáková K. & Hejkal J. 2024 & Hejkal J. 2024. New records of some insects from Strandzha Mts and adjacent Black Sea coast. *ZooNones*, Supplement 16: 1-20.
- Emeljanov A. F. 1974. Proposal on the classification and nomenclature of areals. *Entomologicheskoe Obozrenie* 53 (3): 497-522. (In Russian with English summary).
- Emeljanov A. F. 2018. Sectorial attribution of the range types proposed by K. B. Gorodkov for northern palaeartic. *Entomologicheskoe Obozrenie* 97 (1): 47-59. (In Russian with English summary).
- Ficetola G., Mazel F. & Thuiller W. 2017. Global determinants of zoogeographical boundaries. *Nature Ecology & Evolution* 1 (0089): 1-7.
- Galabov Zh. 1966. Staroplaninska verzihna sistema. In: Gerasimov I. & Galabov Zh. (Eds): *Geography of Bulgaria*. 1. Sofia, Bulgarian Academy of Sciences, 80-116. (In Bulgarian).
- Georgiev B. & Nikolov P. 2010. Invertebrates in the Pomorie Wetland. In: Collection with reports Pomorie Lake region (Protected Site "Pomorie Lake", Ramsar Site, NATURA 2000 sites "Pomorie" BG 0000620 and "Pomorie Lake" BG 0000152), 72-87. (In Bulgarian).
- Georgiev G. & Hubenov Z. 2006. Vertical distribution and zoogeographical characteristics of Cerambycidae (Coleoptera) family in Bulgaria. *Acta Zoologica Bulgarica* 58 (3): 315-343.
- Georgiev V. 1978. Problems of zoogeographical regional division in Bulgaria. *Acta Zoologica Bulgarica* 11: 41-48.
- Georgiev V. 2002. Zoogeographic regioning. In: Copralev I. (Ed.). *Geography of Bulgaria*. Sofia, ForCom, 373-375. (In Bulgarian with English summary).
- Geptner V. 1936. *Obshtaya zoogeografiya*. Moscow, Biomedgiz. 548 pp. (In Russian).
- Gorodkov K. B. 1984. Ranges types of insects of tundra and forests zones of European Part of USSR. In: *Provisional Atlas of the insects of the European Part of USSR*. Leningrad, Nauka, 3-20. (In Russian).
- Grehan J. 1988. The natural biogeographic regions. *Revista di Biologia. Biology Forum* 81 (4): 569-575.
- Grehan J. 1993. Conservation biogeography and the biodiversity crisis: a global problem in space/time. *Biodiversity Letters* 1: 134-140.
- Gruev B. 1988. *General biogeography*. Sofia, Nauka i iskustvo. 396 p. (In Bulgarian)
- Gruev B. 1995. About the mediterranean faunistic complex in Bulgaria. *Annuaire de l'Université de Sofia "Kliment Ohridski", Faculté de Biologie* 86/87 (1- Zoology): 75-82.
- Gruev B. 2000a. Souteuropean peninsular endemism of Alticinae (Coleoptera: Chrysomelidae). *Travaux Scientifiques Université de Plovdiv, Animalia*, 36 (6): 35-50.
- Gruev B. 2000b. About the Atlantic faunistic element in Bulgaria. *Travaux Scientifiques Université de Plovdiv, Animalia*, 36 (6): 67-72. (In Bulgarian, English summary)
- Gruev B. 2000c. About the submediterranean zone of the Palaeartic realm and the submediterranean faunistic element in Bulgaria. *Travaux Scientifiques Université de Plovdiv, Animalia* 36 (6): 73-94. (In Bulgarian with English summary).
- Gruev B. 2002. A comparative study on Alticinae (Coleoptera: Chrysomelidae) in the Balkan Peninsula and Asiatic Turkey. Causes of the similarities and the differences of the fauna. *Travaux Scientifiques Université de Plovdiv, Animalia*, 38(6): 49-79.
- Gruev B. & Bechev D. 2000. Zoogeographical belonging of the leaf beetle species of Lamprosomatinae, Eumolpinae, Chrysomelinae, Alticinae, Hispinae and Cassidinae (Coleoptera, Chrysomelidae) and their distribution in the biogeographical regions of Bulgaria. *Travaux Scientifiques Université de Plovdiv, Animalia* 36 (6): 5-34. (In Bulgarian with English summary).
- Gruev B. & Kusmanov B. 1994. *General biogeography*. Sofia,

- Sofia University K. Ohridski. 498 p. (In Bulgarian).
- Gruev B. & Kusmanov B. 1999. General biogeography. Plovdiv, Plovdiv University. 344 p. (In Bulgarian).
- Heiss E. & Josifov M. 1990. Vergleichende Untersuchung über Artenspektrum, Zoogeographie und Ökologie der Heteropteren-Fauna in Hochgebirgen Österreichs und Bulgariens. Ber. Naturwiss.-med. Vereins Innsbruck, 77: 123-161.
- Holt B., Lessard J.-P., Borregaard M., Fritz S., Araújo M., Dimitrov D., Fabre P.-H., Graham C., Graves G., Jønsson K., Nogués-Bravo D., Wang Z., Whittaker R., Fjeldså J. & Rahbek C. 2013. An Update of Wallace's Zoogeographic Regions of the World. *Science* 339: 74-78.
- Hradský M. & Moucha J. 1964. Raubfliegen (Diptera, Asilidae) Bulgariens. *Acta Faunistica Entomologica Musei Nationalis Pragae* 10 (89): 23-30.
- Hradský M. & Moucha J. 1967. Raubfliegen-Fauna (Diptera, Asilidae) von Bulgarien. *Fragmenta Faunistica* 13 (18): 315-323.
- Hristova N. 2012. Hydrology of Bulgaria. Sofia, Tip-Top, 830 pp. (In Bulgarian).
- Hubenov Z. 1996. Faunistic diversity of Bulgaria – Invertebrates. *Historia Naturalis Bulgarica* 6: 11-16. (In Bulgarian, English summary).
- Hubenov Z. 2008a. Recent fauna of Bulgaria – Animalia: Invertebrata. *Acta Zoologica Bulgarica* 60 (1): 3-21.
- Hubenov Z. 2015. Areographical structure of the Bulgarian non-marine invertebrate fauna (Metazoa: Invertebrata). *Acta Zoologica Bulgarica* 67 (2): 203-213.
- Hubenov Z. 2017. Vertical distribution and comparative zoogeographical characteristic of dipteran fauna (Insecta: Diptera) according to the vegetation belts of the Pirin and Rila Mountains. *Historia Naturalis Bulgarica* 24: 61-119.
- Hubenov Z. 2019. Distribution and Comparative Characteristics of the Dipteran Fauna (Insecta: Diptera) of the Vrachanska Planina, Vitosha, Rila and Pirin Mountains, Bulgaria. *Acta Zoologica Bulgarica* 71 (4): 2019: 501-518.
- Hubenov Z. 2021a. Species composition and distribution of the dipterans (Insecta: Diptera) in Bulgaria. Sofia, Pensoft and National Museum of Natural History (Bulgarian Academy of Sciences), 276 pp. <https://doi.org/10.3897/ab.e68616>
- Hubenov Z. 2021b. Zoogeographical zoning of Bulgaria. *Priroda* 3: 62-70. (In Bulgarian).
- Hubenov Z. 2025a. The Dipterans (Insecta: Diptera) of the Stara Planina Mountains. *Acta Zoologica Bulgarica* 77 (1): 23-39.
- Hubenov Z. 2025b. Species composition and distribution of the dipterans (Insecta: Diptera) of the High Bulgarian Mountains. *Acta Zoologica Bulgarica* 77 (2): 149-172.
- Hubenov Z, Kenderov L. & Pandourski I. 2015. Invertebrate Animals (Metazoa: Invertebrata) of the Atanasovsko Lake, Bulgaria. *Historia Naturalis Bulgarica* 22: 45-71.
- Jacentkovsky D. 1936. Beitrag zur Kenntnis der Raupenfliegen (Tachinariae, Diptera) Bulgariens. *Mitteilungen aus dem Konigl. Naturwissenschaftlichen Instituten in Sofia* 9: 109-134.
- Jacentkovsky D. 1937. Příspěvek k studiu biologické obrany proti lesním škůdcům. *Sborník vysoké školy zemědělské v Brně, ČSR., Fakulta Lesnická, D24*, 54 p.
- Ježek J., Manko P. & Oboňa J. 2020. Synopsis of the Psychodidae (Diptera) fauna of Bulgaria. *Zootaxa* 4877 (2): 201-240. <https://doi.org/10.11646/zootaxa.4877.2.1>
- Josifov M. 1981. Nasekomite ot razred Heteroptera na Balkanskiya poluostrrov. Dissertation (Inst. Zool.), Sofia, Bulg. Acad. Sci., 31-288. (In Bulgarian).
- Josifov M. 1986. Verzeichnis der von der Balkanhalbinsel bekannten Heteropterenarten (Insecta, Heteroptera). *Faun. Abh. Mus. Tierk. Dresden*, 14(6): 61-93.
- Josifov M. 1988. Über den zoogeographischen Charakter der Südeuropäischen Insektenfauna unter besonderer Berücksichtigung der Heteropteren. *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 75: 177-184.
- Josifov M. 1999. Heteropterous insects in the Sandanski-Petrich Kettle, Southwestern Bulgaria. *Historia Naturalis Bulgarica*, 10: 35-66.
- Kechev M. 2021. Data on the distribution of Dolichopodidae (Diptera: Empidoidea) in Bulgaria, with first records for the country. *Historia Naturalis Bulgarica* 42: 95-102.
- Kechev M., Naglis S., Tonguç & Pollet M. 2020. Checklist of the Dolichopodidae (Diptera, Empidoidea) of the Balkan Peninsula, with first records for Bulgaria, Montenegro, North Macedonia, and for the European part of Turkey. *Zootaxa* 4819 (3): 436-472.
- Kovachev S., Stoichev S. & Hainadjieva V. 1999. The zoobenthos of several lakes along the Northern Bulgarian Black Sea coast. *Lauterbornia* 35: 33-38.
- Krastev T. 2002. Morphosculpture of the Bulgarian Black Sea Coast. In: Copralev I. (Ed.). *Geography of Bulgaria*. Sofia, ForCom, 57-60. (In Bulgarian with English summary).
- Krzemiński W. 1984. Limoniidae of Bulgaria (Diptera, Nematocera). I. *Acta Zoologica Bulgarica* 24: 27-34.
- Krzemiński W. & Starý J. 1989. Limoniidae of Bulgaria (Diptera, Nematocera). II. *Bulletin Entomologique de Pologne* 59 (1): 253-279.
- Kryzhanovsky O. 1965. Composition and origin of the terrestrial fauna of Central Asia (based chiefly on the beetles material). Moscow-Leningrad, Nauka. 430 pp. (In Russian).
- Kryzhanovsky O. 1976. On the principles of zoogeographical regionalization of land. *Zoologicheskyy Zhurnal* 55 (7): 965-975. (In Russian with English summary).
- Kryzhanovsky O. 2002. Composition and distribution of entomofauna of the globe. Moscow, KMK. 237 pp. (In Russian).
- Lattin de G. 1967. *Grundriss der Zoogeographie*. Hochschulverlag für Biologie. 12. Jena, VEB Gustav Fischer Verlag. 602 pp.
- Lavčiev V. 1969. Forschungen über die echten Fliegen (Muscidae, Diptera) im Reservat Ropotamo. *Bulletin de l'Institut de Zoologie et Musée* 30: 59-70. (In Bulgarian, German summary).
- Lavčiev V. 1980. Muhite okolo nas i borbata s tyah [The flies around us and the fight against them]. Sofia, Bulgarian Academy of Sciences. 205 pp. (In Bulgarian).
- Lavčiev V. 2003. *Catalogus faunae bulgaricae*. 5. Diptera: Fannidae, Muscidae, Stomoxydidae. Sofia, Editio Academica "Professor Marin Drinov". 78 pp.
- Lopatín I. K. 1989. Zoogeography. Minsk, Vysheyschaya shkola. 392 pp. (In Russian).
- Löw H. 1862. Über einige bei Varna gefangene Dipteren. *Wiener Entomologische Monatschrift* 6 (6): 161-175.
- Lutovinova E., Kanavalova L., Barták M. & Kokan B. 2022. New or rare records for the Balkan Tachinidae (Diptera: Oestroidea). *Acta Zoologica Bulgarica* 74 (2): 175-182.
- Malicky H., Ant H., Aspöck H., de Jong R., Thaler K. & Varga Z. 1983. Argumente zur Existenz und Chorologie mit-

- teleuropäischer (extramediterranean-europäischer) Faunen-Elemente. *Entomologia Generalis* 9: 101-119.
- McAlpine J. F. 1989. Phylogeny and classification of the Muscomorpha. In: McAlpine J. F. & Wood D. M. (Eds.). *Manual of Nearctic Diptera* 3. Ottawa, Research Branch Agriculture Canada. pp. 1397-1518.
- Mendl H. 1986. Stelmücken (Limoniidae) aus Bulgarien (Diptera, Nematocera). *Spixiana* 9 (3): 275-282.
- Michailova P. 1973a. Triploidy in *Clunio marinus* Haliday from the Bulgarian Black Sea coast. *Comptes Rendus de l'Académie Bulgare des Sciences* 26 (11) : 1541-1543.
- Michailova P. 1973b. Untersuchungen über den Chromosomenpolymorphismus bei *Chironomus salinarius* Kieff., *Chironomus valkanovi* Michailova und *Chironomus anchialicus* Michailova (Diptera, Chironomidae) von der bulgarischen Schwarzmeerküste. *Zoologischer Anzeiger* 191 (5/6): 348-364.
- Michailova P. 1974. Zwei neue Arten der Gattung *Chironomus* (Diptera, Chironomidae) von der bulgarischen Schwarzmeerküste. *Zoologische Beiträge* 20 (2): 339-357.
- Michailova P. 1980. A review of the European species of genus *Clunio* Haliday, 1855 (Diptera, Chironomidae). *Zoologischer Anzeiger* 205 (5/6): 417-432.
- Michailova P. 1982. Chironomids (Diptera, Chironomidae) in fish raising pools in Bulgaria. *Acta Zoologica Bulgarica* 20: 51-68. (In Bulgarian with English summary).
- Michailova P. 1989. The polytene chromosomes and their significance to the systematics and phylogeny of the family Chironomidae, Diptera. *Acta Zoologica Fenica* 186: 107 pp.
- Mikov O. 2005. History of investigations of malaria mosquitoes in Bulgaria. *Infectology* 42 (4): 39-47. (In Bulgarian).
- Mikov O. 2008. Biologichni, ekologichni i epidemiologichni prouchvaniya na anofeliyni komari v nyakoi regioni na Bulgariya [Biological, ecological and epidemiological studies of the anopheline mosquitoes in some regions of Bulgaria]. Ph.D. thesis, Sofia, National Center of Infectious and Parasitic Diseases. 155 pp. (In Bulgarian).
- Mikov O., Nikolov G., Schaffner F. & Mathis A. 2013. First record and establishment of *Aedes albopictus* in Bulgaria. In: VBORNET-EMCA Joint Meeting "Invasive mosquitoes and public health in the European context", Antwerp, Belgium, 28-29 November 2013 (Conference paper).
- Müller P. 1974. Aspects of zoogeography. The Hague, W. Junk Publishers. 208 pp.
- Müller P. 1980. Biogeographie. Stuttgart, Ulmer. 414 pp.
- Nagatomi. A. 1996. An essay on phylogeny of the orthorrhaphous Brachycera (Diptera). *Entomologist's Monthly Magazine* 132: 95-148.
- Nartshuk E. P. 2003. Key to families of Diptera (Insecta) of the fauna of Russian and adjacent countries. St. Petersburg, Proceedings of the Zoological Institute 294, Russian Academy of Sciences. 251 pp. (In Russian).
- Nedelkov N. 1912. Sixth contribution to the entomological fauna of Bulgaria. *Revue Academy Sciences de Bulgarie* 2: 177-218. (In Bulgarian).
- Oosterbroek P. 2006. The European families of the Diptera. Identification, diagnosis, biology. KNNV Publishing. 209 pp. DOI: 10.1163/9789004278066
- Pape T., Blagoderov V. & Mostovski M. 2011. Order Diptera Linnaeus, 1758. In: Zhang Z.-Q. (Ed.). *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa* 3148: 222-229.
- Papp L. & Darvas B. (Eds.). 1997. *Contributions to a Manual of Palaearctic Diptera* 2. Nematocera and Lower Brachycera. Budapest, Science Herald. 592 pp.
- Papp L. & Darvas B. (Eds.). 1998. *Contributions to a Manual of Palaearctic Diptera* 3. Higher Brachycera. Budapest, Science Herald. 880 pp.
- Papp L. & Darvas B. (Eds.). 2000a. *Contributions to a Manual of Palaearctic Diptera* 4: Appendix. Budapest, Science Herald. 604 pp.
- Papp L. & Darvas B. (Eds.). 2000b. *Contributions to a Manual of Palaearctic Diptera* 1: General and Applied Dipterology. Budapest, Science Herald. 978 pp.
- Popov A. 2002. Zoogeographical analysis of Neuroptera in Bulgaria. *Acta Zoologica Hungarica* 48 (Suppl. 2): 271-180.
- Povolný D. & Verves Y. 1990. A preliminary list of Bulgarian Sarcophaginae (Diptera). *Acta Entomologica Musei Nationalis Pragae* 43: 283-329.
- Procheş Ş. & Ramdhani S. 2012. The World's zoogeographical regions confirmed by cross-taxon analyses. *BioScience* 62 (3): 260-270.
- Sabev L. & Stanev S. 1959. Klimatichnite rayoni na Balgaria i tehniyat klimat. Proceedings of the Institute of Hydrology and Meteorology 5: 29-174. (In Bulgarian).
- Skuhrava M., Skuhravy V., Dončev K. & Dimitrova B. 1991. Gall midges (Cecidomyiidae, Diptera) of Bulgaria. I. Faunistic researches in the 1978-1987 period. *Acta Zoologica Bulgarica* 42: 3-26.
- Skuhrava M., Skuhravy V., Dončev K. & Dimitrova B. 1992. Gall midges (Cecidomyiidae, Diptera) of Bulgaria. II. Host plant relations and economic importance. *Acta Zoologica Bulgarica* 43: 23-42.
- Slípka J. 1959. *Symplectomorpha stictica* Meig. (Diptera, Nematocera) als fakultativer Halobiont. *Bulletin de l'Institut de Zoologie et Musee (Academie Bulgare des sciences)* 8: 105-108.
- Soós A. & Pap L. (Eds.). 1984-1993. *Catalogue of Palaearctic Diptera* 1-13. Budapest, Hungarian Natural History Museum.
- Sørensen T. 1948. A method of establishing groups of equal amplitude in plant sociology based on similarity of species content. *Kongelige Danske Videnskabernes Selskab. Biologiske Skrifter* 4: 1-34.
- Stanev S. 1991. Climatic regions of Bulgaria. In: Stanev S., Kyuchukova M. & Lingova St. (Eds.), *The climate of Bulgaria*. Sofia, Bulgarian Academy of Sciences, 53-89. (In Bulgarian with English summary).
- Starý J. 1982. New species of Limoniidae from the Mediterranean (Diptera). *Bulletin Zoologisch Museum Universiteit van Amsterdam* 9: 25-30.
- Starý J. 2009. The identity of *Dicranomyia (Dicranomyia) luteipennis* Goetghebuer (Diptera, Limoniidae). *Zootaxa* 2155: 55-68.
- Starý J. & Krzemiński W. 1993. Additions to the list of Bulgarian Limoniidae and Pediciidae (Diptera). *Acta Zoologica Cracoviensia* 35: 569-572.
- Stoichev S. 1996. On the Chironomid fauna from Bulgarian inland waters. *Lauterbornia* 25: 117-123.
- Stoichev S. 1998. The zoobenthos from the lakes Schabla-Ezerets (northern Black Sea coast of Bulgaria). In: Golemansky V., Naidenov W. (Eds.). *Biodiversity of Schabla Lake System, Sofia, "Prof. Marin Drinov" Academic Publishing House*. pp. 91-99.

- Stojanov N. 1966. Rastitelna pokrivka. In: Beshkov A. (Ed.). Geography of Bulgaria. 1. Sofia, Bulgarian Academy of Sciences, 447-482. (In Bulgarian).
- Szilády Z. 1934. Beiträge zur Dipterenfauna Bulgariens. Bulletin de la Société Entomologique de Bulgarie 8: 145-151.
- Tichkov H. 1976. Le climat des régions montagneuses de la Bulgarie. Structure et genèse. Sofia, Bulgarian Academy of Sciences, 210 pp. (In Bulgarian with French summary).
- Todorova E. 2014. Kachestven i kolichestven sastav na nyakoi osnovni grupi bezgrabnachni zhivotni s vazhno znachenie za ezeroto Vaya. [Qualitative and quantitative composition of some main groups of invertebrates of importance for Vaya Lake]. Ph.D. thesis, Sofia University, Faculty of Biology (Department Zoology and Anthropology). 276 pp. (In Bulgarian).
- Udvardi M. 1975. A classification of the Biogeographical Provinces of the World. Morges, IUCN Paper 18. 49 pp.
- Valkanov A. 1936. Notizen über die Brackwässer Bulgariens. II. Versuch einer hydrographischen und biologischen Erforschung derselben. Annuaire de l'Université de Sofia. Faculté physico-mathématique 32 (3): 209-341. (In Bulgarian with German summary).
- Valkanov A. 1948. *Thalassomyia frauenfeldi* Schiner vom Schwarzen Meer. Arbeiten aus der Biologischen Meeresstation in Varna, 14: 103-112. (In Bulgarian with German summary).
- Valkanov A. 1954. Betrag zur Kenntnis unserer Schwarzmeerfauna. Arbeiten aus der Biologischen Meeresstation in Varna, 18: 49-53. (In Bulgarian with German summary).
- Valkanov A. 1957. Katalog unserer Schwarzmeerfauna. Arbeiten aus der Biologischen Meeresstation in Varna, 19: 1-62. (In Bulgarian with German summary).
- Varadinova E., Vassilev V., Stoichev S. & Uzunov Y. 2010. The macrozoobenthos of the Pomorie Lake. In: Collection with reports Pomorie Lake region (Protected Site „Pomorie Lake”, Ramsar Site, NATURA 2000 sites „Pomorie” BG 0000620 and „Pomorie Lake” BG 0000152), 95-100. (In Bulgarian).
- Varbanov M. 2002. Lakes and swamps. In: Geography of Bulgaria. Sofia, ForKom: 237-242. (In Bulgarian).
- Vassilev V., Konsulov A. 1998. Zooplankton in the lake Pomoriysko – composition, dynamics, trophic interactions and secondary assimilation. Proceedings of the Institute of Oceanology (Varna), 2, 146-153. (In Bulgarian with English summary).
- Velchev V. 2002. Characteristic features and regularities in the distribution of the present-day vegetation. In: Copralev I. (Ed.). Geography of Bulgaria. Sofia, ForKom, 321-324. (In Bulgarian with English summary).
- Velchev V., Ganchev S. & Bondev I. 1982. Rastitelni poyasi. In: Galabov Zh. (Ed.). Geography of Bulgaria. 1. Sofia, Bulgarian Academy of Sciences, 439-443. (In Bulgarian with English summary).
- Velev St. 2002. Climatic regioning. In: Copralev I. (Ed.). Geography of Bulgaria. Sofia, ForKom, pp. 155-156. (in Bulgarian with English summary).
- Velev St. 2010. Climate districts and regions. In: Copralev I. (Ed.). Bulgaria geographic atlas. Sofia, Tangra, p. 89. (In Bulgarian with English summary).
- Vigna Taglianti A., Audisio P., Biondi M., Bologna M., Carpaneto G., Biase A., Fattorini S., Piattella E., Sindaco R. 1999. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic Region. Biogeographia 20: 31-59.
- Wagner R. & Joost W. 1988. Psychodidae aus Bulgarien (Insecta, Diptera). Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden B, 16 (3): 29-34.
- Wood D. M. & Borkent A. 1989. Phylogeny and classification of the Nematocera. In: McAlpine J. F. & Wood D. M. (Eds.). Manual of Nearctic Diptera 3. Ottawa, Research Branch Agriculture Canada. pp. 1333-1370.
- Woodley N. E. 1989. Phylogeny and classification of the “Orthorhaphous” Brachycera In: McAlpine J. F. & Wood D. M. (Eds.). Manual of Nearctic Diptera Volume 3. Ottawa, Research Branch Agriculture Canada. 1371-1395.
- Yeates D. & Wiegmann B. 1999. Congruence and Controversy: Toward a Higher-Level Phylogeny of Diptera. Annual Review of Entomology 44 (1):397-428. DOI: 10.1146/annurev.ento.44.1.397 Source: PubMed
- Yeates D., Wiegmann B., Courtney G., Meier R., Lambkin Ch. & Pape T. 2007. Phylogeny and systematics of Diptera: Two decades of progress and prospects. Zootaxa 1668: 565-590.
- Zilahi G. 1934. Beiträge zur Fliegenfauna Bulgariens. I. Chironomidae. Mitteilungen der Bulgarischen Entomologischen Gesellschaft in Sofia 8: 152-158.
- Ziegler J. 2003. Ordnung Diptera, Zweiflügler (Fliegen und Mücken). In: Dathe H. (Ed.). Lehrbuch der Speziellen Zoologie, Band I: Wirbellose Tiere, 5. Teil: Insecta. Heidelberg, Spektrum. pp. 756-860.
- Zielke E. 2016a. Update of distribution records of *Phaonia* Robineau-Desvoidy (Diptera: Muscidae) from Bulgaria with the description of a new species. Journal of Entomology and Zoology Studies 4 (4): 626-632.
- Zielke E. 2016b. Update of distribution records of Mydaeinae (Diptera: Muscidae) from Bulgaria. Journal of Entomology and Zoology Studies 4 (6): 310-315.
- Zielke E. 2018. An update of distribution records of *Helina* R.-D. and other genera of the subfamily Phaoniinae (Diptera: Muscidae) from Bulgaria. Acta Musei Moraviae, Scientiae biologicae 103 (2): 249-267.
- Zielke E. 2019. Update of distribution records of *Thricops* Rondani and other genera of the subfamily Azeliinae (Diptera: Muscidae) from Bulgaria. Journal of Entomology and Zoology Studies 7 (3): 316-323.

Received: 05.09.2025

Accepted: 28.01.2026