

Fish Community in Karla Reservoir (Central Greece) after the Filling Stage

Maria Chamoglou^{1*} & Apostolos Apostolou²

¹ Management Body of Eco-development Area of Karla – Mavrovouni – Kefalovriso – Velestino, 38500 Kanalia, Greece

² Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 2 Gagarin Street, 1113 Sofia, Bulgaria

Abstract: Karla Lake was drained in 1962 and restored as a reservoir in 2009. This re-established water body is considered a vital aquatic ecosystem as it is recognised as a Special Area of Conservation under the EC Habitat Directive (92/43/EEC). As recommended by the European Water Framework Directive, in 2013 and 2016 the fish community was characterised. A total of 8,974 specimens weighting 130.6 kg and belonging to 11 species and six families (Centrarchidae, Clupeidae, Cobitidae, Cyprinidae, Gobiidae and Poeciliidae) were caught. Four species (*Alburnus thessalicus*, *Vimba melanops*, *Squalius vardarensis* and *Knipowitschia thessala*) are considered as endemic to Greece and the Balkan Peninsula and three (*Lepomis gibbosus*, *Carassius gibelio*, *Gambusia holbrooki*) are alien. The reservoir fish community was dominated by species of the family Cyprinidae, both in terms of number and weight, supporting the eutrophic character of the water body. NPUE and BPUE values did not differ significantly between the two sampling campaigns (Wilcoxon's test, $P > 0.05$). The current results could be used as a database for future comparisons and for management and conservation of fish assemblages. One sampling every three years (as required by the European Water Framework Directive) is assumed as insufficient in such rapidly developing ecosystems.

Key words: fish monitoring, fish community, fish assessment, lake restoration

Introduction

Recently, restoration of lakes presents a significant part of aquatic ecosystems' management. Restoration activities have been taken all over the world, including Europe. In most cases, the actions taken concern the balance of material and/or certain ecosystem components e.g. macrophytes (HILTA et al. 2006). In Greece restoration measures have been undertaken in Lake Pamvotis (MYLOPOULOS et al. 2007) but the restoration of a drained lake as Karla represents a pilot challenge for the country.

Karla Lake was drained in 1962 and rebuilt as a reservoir in 2009. This newly re-established water body is considered a vital aquatic ecosystem as it is recognised as a Special Area of Conservation (SAC) under the EC Habitats and Species Directive (92/43/EEC). Restoration activities as construction of dykes

and floodgates were undertaken in 2000, following the Ramsar guidelines (ZALIDIS et al. 2004).

The reservoir inflows derive primarily by the diversion of the Pinios River floods, the surface runoff of the watershed and the direct precipitation on the reservoir surface. The annual water abstraction from the Pinios River to the reservoir is 80 km³ for the period mid-October to mid-April each year (MINISTRY OF ENVIRONMENT, PLANNING AND PUBLIC WORKS 2000).

At present, the reservoir suffers from a large-scale water deviation. Since September 2009 (start of refilling) it has not still achieved the lowest ecological limit that would support the functions of the wetland, as set by the environmental terms of the project (at +46.4 m) (SIDIROPOULOS et al. 2015). Moreover,

*Corresponding author: mchamoglou@gmail.com

cyanobacterial water blooms during the warm period have been reported (OIKONOMOU et al. 2012, NIKOULI et al. 2013). The reservoir has been designated as a nature wildlife shelter and as part of the NATURA 2000 network (code GR 1420004 “Karla - Maurovouni - KefalovrisoVelestino - Neohori”).

In the past, before its drying in 1962, the Karla Lake was among the most important lakes in Greece as far as fisheries were concerned. It was a place of cultural traditions and development of an impressively unique lifestyle of the people who – for centuries – have been living off fishing (ROUSKAS 2001). The

only scientific data for the fish fauna of the former lake derive from a single survey. According to ANANIADIS (1956), fish fauna of the Karla Lake was characterised by the family of Cyprinidae. The most abundant species were *Cyprinus carpio* Linnaeus, 1758 along with *Rutilus rutilus* (Linnaeus, 1758) and *Scardinius erythrophthalmus* (Linnaeus, 1758). The population of *Alburnus thessalicus* Stephanidis, 1950 was declining in terms of abundance, while *Anquilla anquilla* (Linnaeus, 1758) was rare (ANANIADIS 1956). The former Karla Lake has remained drained for 47 years until the recent construction of the reservoir. Any




Fig. 1. The studied area of the Karla Reservoir with the sampling stations.

scientific information for the fish community of the drainage (channels and 12 small reservoirs) except a technical report (ECONOMIDIS et al. 2004) is lacking.

At present, in the artificial ecosystem the fishing is prohibited by Greek law.

The objective of the study was to clarify the fluctuations in the structure and abundance of the newly-formed fish community after the restoration of a drained lake (Karla).

Materials and Methods

Study area

Karla Reservoir (39°29'02''N, 22°51'41''E) is located in the south-east part of Thessaly Region, Central Greece. It is constructed as part of the project 'Recreation of Lake Karla' funded initially by the 3rd CSF 2000-2006 and subsequently by the NSRF 2007-2013. According to WFD, the Karla

Reservoir, in terms of typology is considered a heavily modified water body (Management Plan of River Basin Thessaly (GR 08)). It represents an eutrophic (CHAMOGLIOU et al. 2014) reservoir at an altitude of 44 m a.s.l, drainage area of 1,171 km², surface area of 38 km² and a maximum depth 4.5 m.

Sampling

Fish sampling was carried out in April 2013 and 2016 with type Nordic benthic gillnets (according to CEN, 2005, Water Quality – Sampling of Fish with Multimesh Gillnets) by performing stratified random sampling (APPELBERG 2000). The fishing effort was set according to the area (38 km²) and the maximum sampling depth (\approx 4.5 m) of the reservoir corresponding to 12 sampling points (Fig. 1). Nets were set before dusk and hauled after dawn, in order to ensure a standard soak time of 12 hours for two days (6 nets on the first day and 6 the other). Nets were

Table 1 Catch per unit effort in terms of (a) number (NPUE, ind./m²) and (b) weight (BPUE, g/m²) of fish species from Karla Reservoir in 2013 and 2016. N – native species, A – alien; EN – Endemic; ENNO – Endemic to Northern Greece, including Thessaly; ENBA – Endemic to the Balkans, MED – Mediterranean; NA – non-applicable; + presence; - absence.

Year	ANANIADIS (1956)	2013		2016		Mean difference in TL (cm)	Mean difference in W (gr)	Origin (BARBERI et al. 2015)	Endemicity (ECONOMOIOU et al. 2007)
		NPUE	BPUE	NPUE	BPUE				
Family/Species								(1)	(2)
Anguillidae									
<i>Anguilla anguilla</i>	+	-	-	-	-	-	-	N	
Centrarchidae									
<i>Lepomis gibbosus</i>	-	1.07	27.12	0.517	5.416	-2.818	-14.08	A	
Clupeidae									
<i>Alosa fallax</i>	-	0.007	3.772	-	-	NA	NA	N	MED
Cobitidae									
<i>Cobitis</i> sp.	-	0.083	0.658	0.006	0.023	0.185	-3.707	N	EN
Cyprinidae									
<i>Alburnus thessalicus</i>	+	3.744	49.876	10.291	81.974	-1.557	-5.681	N	ENBA
<i>Carassius gibelio</i>	-	0.074	6.056	0.026	5.338	5.341	124.151	A	
<i>Cyprinus carpio</i>	+	0.17	8.602	0.165	24.568	4.917	98.574	N	
<i>Rutilus rutilus</i>	+	0.296	26.596	0.044	1.001	-6.12	-63.749	N	
<i>Scardinius erythrophthalmus</i>	+	-	-	-	-	-	-	N	
<i>Squalius vardarensis</i>	-	-	-	0.013	0.161	NA	NA	N	ENBA
<i>Vimba melanops</i>	-	-	-	0.024	0.722	NA	NA	N	ENBA
Gobiidae									
<i>Gobio feraeensis</i>	+	-	-	-	-	-	-	N	ENNO
<i>Knipowitschia thessala</i>	-	0.002	0.005	0.065	0.051	3.857	0.789	N	ENNO
Poeciliidae									
<i>Gambusia holbrooki</i>	-	0.02	0.044	-	-	NA	NA	A	




Fig. 2. Percentage (%) participation of fish species by family in terms of a) abundance and b) weight caught at the Karla Reservoir in 2013 (blue colour) and 2016 (red colour). Clupeidae is not included, since only four specimens were captured in 2013.

set in the same location in both surveys following the same methodology. A handheld GPS was used to mark the precise location of each net.

Fish samples were identified to the species level according to KOTTELAT & FREYHOF (2007), with the exception of species of Cobitidae that were identified at the generic level (*Cobitis* sp.). Some of the caught specimens expressed specific external characteristics, such as *C. stephanidisi* Economidis, 1992, others, such as *C. vardarensis* Karaman, 1928 showed intermediate characteristics. Due to the fact that hybridisation in the genus *Cobitis* is common (CHOLEVA et al., 2008), it is safer to identify these samples to genus level, until molecular-genetic analyses enlighten this issue.

Statistical analysis

Catches were analysed separately for each survey and the total number and weight of specimens per species were recorded. For each species, CPUE was calculated as the number of individuals per m² net (NPUE) and the weight of fish per m² net (BPUE; APPELBERG 2000). The values of CPUE per survey were compared by Wilcoxon's test. Mean difference in TL (cm) between the two samplings (mean TL during second - mean TL during first sampling) were also calculated.

Results

These are the first data on the fish fauna of the Karla Reservoir. Based on the results of both campaigns, a total of 8974 specimens weighting 130.6 kg of 11 fish species (Table 1) and six families (Centrarchidae, Clupeidae, Cobitidae, Cyprinidae, Gobiidae, Poeciliidae) were identified. Eight are native, whereas *Lepomis gibbosus* (Linnaeus, 1758), *Carassius gibelio* (Bloch, 1782) and *Gambusia holbrooki* Girard, 1859 are alien (BARBIERI et al. 2015). Among the native species, five species are considered as endemic to Greece and the Balkan Peninsula

(ECONOMOU et al. 2007). *Alosa fallax* (Lacepède, 1803) is listed in annexes II and V of the EU Habitats and Species Directive. Both sampling campaigns revealed a fish community dominated by cyprinids (6 out of 11 species).

Comparing the samples collected in 2016 with those taken in 2013, the fish fauna of the Karla Reservoir was similar in terms of species composition. However, we recorded between-year differences in abundance for some fish families. Cyprinidae increased by 16.2% in terms of abundance (Fig. 2). The second most abundant family Centrarchidae, represented by *L. gibbosus*, showed a decrease of 15% from 2013 to 2016. The decline in abundance of *L. gibbosus* and *G. holdbrooki*, should be probably considered as positive for native fish (MILLS et al. 2004, COPP et al. 2005).

Comparison of the overall sampling results did not show significant differences (Wilcoxon test values for NPUE and BPUE were 33 and 28, respectively, at $P > 0.05$). Moreover, samplings were performed when water level was similar. However, the examination of the values of NPUE and BPUE that accounted for each species revealed some differences between the surveys. A considerable increase in terms of NPUE was observed for *K. thessala* from 0.002 individuals/m² in 2013 to 0.065 ind./m² in 2016, followed by *A. thessalicus* from 3.744 ind./m² in 2013 to 10.291 ind./m² in 2016. On the contrary, a decline in terms of NPUE was displayed by *Cobitis* sp. (from 0.083 ind./m² in 2013 to 0.006 ind./m² in 2016), *R. rutilus* (from 0.296 ind./m² in 2013 to 0.044 ind./m² in 2016), *C. gibelio* (from 0.074 ind./m² in 2013 to 0.026 ind./m² in 2016) and *L. gibbosus* (from 1.07 ind./m² in 2013 to 0.517 ind./m² in 2016; Table 1). The species that increased their biomass were *K. thessala* (from 0.005 g/m² in 2013 to 0.051 g/m² in 2016), *C. carpio* (from 8.602 g/m² in 2013 to 24.568 g/m² in 2016) and *A. thessalicus* (from 49.876 in 2013 g/m² to 81.974 g/m² in 2016).

Discussion

Generally, fish species of the family Cyprinidae are well adapted to eutrophic conditions (PERSSON 1991, MEHNER et al. 2005) similar to those in the Karla Reservoir (CHAMOGLIOU et al. 2013). The other five families that we recorded were represented by a single species. Two cyprinid species, the eurytopic *Vimba melanops* (Heckel, 1837) and *Squalius vardarensis* were recorded only in 2016 and they probably have entered from the Pinios River, where they are abundant (BARBIERI et al. 2015). *Alosa fallax* was recorded for the first time in the reservoir but only in 2013. It has likely entered the reservoir from the Pinios River, most probably during the spawning period. The alien *Lepomis gibbosus* was registered for the first time in the adjacent Pinios River in 2011 (KOUTSIKOS et al. 2012), indicating its further spreading.

Anguilla anguilla, *Gobio feraeensis* and *Scardinius erythrophthalmus*, previously reported by ANANIADIS (1956) for the former Lake Karla, were not detected during our study.

The fact that *G. holbrooki* was present only in 2013 survey needs further investigation, in order to clarify the reasons for the decline of this eurybiont and invasive species.

Karla Reservoir will likely face numerous challenges and changes, regarding the structure of the local fish fauna, since it is still in its initial stage of filling with water from the Pinios River. These challenges are related with the multipurpose use of the reservoir as it has to satisfy both environmental (ecological restoration) and social needs (irrigation, flood protection, water supply of the nearby Volos City and ecotourism activities) in the near future. Furthermore, several variables, such as physical, chemical and biological properties of the water, surface area, age, hydraulic retention time of the reservoir, the composition of the fish fauna in the river and watershed area may influence the composition of the fish community in the reservoir (AGOSTINHO et al. 1999, 2016).

Our results provide initial information on the fish community structure of the Karla Reservoir after its refilling and could be used as a basis for future management, as well as conservation of local fish assemblages. The followed sampling strategy recommended by the WFD 2000/60/EC represents a useful tool for interpreting differences in fish communities based on catch per unit effort (EMMRICH et al. 2011, BRUCET et al. 2013), however, is insufficient in such ecosystems. A sampling every three years is not enough to reflect quick changes, characterising

an ecosystem under restoration, as is the case of the Karla Reservoir. Combining sampling methods is also recommended, in order to explore in detail fish biodiversity and abundance (HUBERT 1996). In the case of the Karla Reservoir, beach seine/long lines/electrofishing could be used altogether with gill nets.

Acknowledgements: This study has been co-funded by (1) The European Union (ERDF: European Regional Development Fund- ERDF) and Greek national funds through the Operational Program “Environment and Sustainable Development” 2007-2013 – Priority Axis “Protecting Nature and Biodiversity”; (2) *Recreation of Lake Karla Project*, funded initially by the 3rd CSF 2000-2006 and subsequently in the NSRF 2007-2013; (3) Water Framework Directive surveillance monitoring program by Management Body of Ecodevelopment Area of Karla – Mavrovouni – Kefalovriso – Velestino, funded by European Union.

References

- AGOSTINHO A. A., MIRANDA L. E., BINI L. M., GOMES L. C., THOMAZ S. M. & SUZUKI H. I. 1999. Patterns of colonization in neotropical reservoirs, and prognoses on aging. In: TUNDISI J. G. & STRAŠKRABA M. (Eds.): Theoretical reservoir ecology and its applications. International Institute of Ecology, São Carlos, pp 585.
- AGOSTINHO A. A., GOMES L. C., SANTOS N. C. L., ORTEGA J. C. G. & PELICICE F. M. 2016. Fish assemblages in Neotropical reservoirs: Colonization patterns, impacts and management. *Fisheries Research* 173: 26–36.
- APPELBERG M. 2000. Swedish standard methods for sampling freshwater fish with multi-mesh gillnets. *Fiskeriverket Information* 2000, 1: 3-32.
- ANANIADIS C. 1956. Limnology study of Lake Karla. *Bulletin de l'Institut Oceanographique* 1083: 1-19.
- BARBIERI R., ZOGARIS S., KALOGIANNI E., STOUMBOUDI M. T., CHATZINIKOLAOU Y., GIAKOUMI S., KAPAKOS Y., KOMMATAS D., KOUTSIKOS N., TACHOS V., VARDAKAS L., ECONOMOU A. N. 2015. Freshwater Fishes and Lampreys of Greece: An annotated checklist. *Monographs on Marine Sciences* No 8. Hellenic Center of Marine Research. Athens, Greece. pp. 130.
- BRUCET S., PÉDRON S., MEHNER T., LAURIDSEN T. L., ARGILLIER C., WINFIELD I. J., VOLTA P., EMMRICH M., HESTHAGEN T., HOLMGREN K., BENEJAM L., KELLY F., KRAUSE T., PALM A., RASK M. & JEPPESEN E. 2013. Fish diversity in European lakes: Geographical factors dominate over anthropogenic pressures. *Freshwater Biology* 58: 1779–1793.
- CEN 2005. Water Quality – Sampling of Fish with Multimesh Gillnets. European Committee for Standardization, EN 14757, Brussels.
- CHAMOGLIOU M., PAPADIMITRIOU T. & KAGALOU I. 2014. Keys-Descriptors for the Functioning of a Mediterranean Reservoir: The Case of a New Lake Karla-Greece. *Environmental processes*, 1:127-135.
- CHOLEVA L., APOSTOLOU A., RAB P. & JANKO K. 2008. Making it on their own: sperm-dependent hybrid fishes (*Cobitis*) switch the sexual hosts and expand beyond the ranges of their original sperm-donors. *Philosophical Transactions of*

- the Royal Society B: Biological Sciences 363: 2911-2919.
- COPP G. H., BIANCO P. G., BOGUTSKAYA N. G., EROS T., FALKA I., FERRIERA M. T., FOX M. G., FREYHOF J., GOZLAN R. E., GRABOWSKA J., KOVAC V., MONERO-AMICH R., NASEKA A. M., PENAZ M., POVZ M., PRYBYLSKI M., ROBILLARD M., RUSSELL I. C., STAKENAS S., SUMER S., VILA-GISPERS A. & WIESNER C. 2005. To be, or not to be, a non-native freshwater fish? *Journal of Applied Ichthyology* 21: 242-262.
- ECONOMIDIS P., BOBORI D. & PERGANTIS F. 2004. Ichthyologic investigation in the area of the former Lake Karla. Final technical report, 157 p. (In Greek).
- ECONOMOU A.N., GIAKOUMI S., VARDAKAS L., BARBIERI R., STOUMBOUDI M. & ZOGARIS S. 2007. The freshwater ichthyofauna of Greece – an update based on a hydrographic basin survey. *Mediterranean Marine Science* 8 (1): 91-166.
- EMMRICH M., BRUCET S., RITTERBUSCH D. & MEHNER T. 2011. Size spectra of lake fish assemblages: Responses along gradients of general environmental factors and intensity of lake-use. *Freshwater Biology* 56: 2316–2333.
- HILTA S., GROSSB E., HUPFERA M., MORSCHIEDC H., MÄHLMANN D J., MELZERE A., POLTZF J., SANDROCKG S., SCHARFG E., SCHNEIDERE S. & VAN DE WEYERH K. 2006. Restoration of submerged vegetation in shallow eutrophic lakes – A guideline and state of the art in Germany. *Limnologica* 36: 155–171.
- HUBERT W. A. 1996. Passive capture techniques. In: MURPHY B. R. & WILLIS D. W. (Eds.): *Fisheries techniques*, 2nd ed. Bethesda: American Fisheries Society, pp. 302.
- KOTTELAT M. & FREYHOF J. 2007. *Handbook of European freshwater fishes*. Cornol, Switzerland pp. 646.
- KOUTSIKOS N. S., ZOGARIS L., VARDAKAS V., TACHOS V., KALOGIANNI E., SANDA R., CHATZINIKOLAOU Y., GIAKOUMI S., ECONOMIDIS P. S. & ECONOMOU A. N. 2012. Recent contributions to the distribution of the freshwater ichthyofauna in Greece, *Mediterranean Marine Science*, 13 (2): 268-277.
- MEHNER T., DIEKMANN M., BRAMICK U. & LEMCKE R. 2005. Composition of fish communities in German lakes as related to lake morphology, trophic state, shore structure and human-use intensity. *Freshwater Biology* 50: 70 – 85.
- MILLS M. D., RADER R. B. & BELK M. C. 2004. Complex interactions between native and invasive fish: the simultaneous effects of multiple negative interactions. *Oecologia* 141: 713-721.
- RIVER BASIN MANAGEMENT PLAN FOR THE WATER DISTRICT OF THESSALY (GR08) 2017. Government Gazette 4682/B'/29-12-2017. Ministry of Environment Energy and Climate Change, Special Secretariat for Water. pp 65997 (In Greek).
- MINISTRY OF ENVIRONMENT, PLANNING AND PUBLIC WORKS 2000. Joint Ministerial Decision 112839/18-12-2000: Modification – Supplementation – Codification of the Environmental Terms for the construction and operation of former lake Karla re-flooding project at Prefectures of Larisa and Magnesia, Athens: Greece pp 17. (In Greek).
- MYLOPOULOS N., MYLOPOULOS Y., KOLOKYTHA E. & TOLIKAS D. 2007. Integrated water management plans for the restoration of lake Koronia, Greece, *Water International*, 32 (Suppl. 1): 720-738.
- NIKOULI E., KORMAS K., PANAGIOTIS B., KARAYANNI H. & MOUSTAKA-GOUNI M. 2013. Harmful and parasitic unicellular eukaryotes persist in a shallow lake under reconstruction (L. Karla, Greece). *Hydrobiologia* 718:73-83.
- OIKONOMOU A., Katsiapi M., KARAYANNI H., MOUSTAKA-GOUNI M. & KORMAS K. 2012. Plankton microorganisms coinciding with two consecutive mass fish kills in a newly reconstructed lake. *The Scientific World Journal*, ID 504135: 14 p.
- PERSSON L. 1991. Interspecific interactions. In I. J. Winfield, & J. S. Nelson (Eds.) *Cyprinid fishes. Systematics, biology and exploitation*. Springer. London 667 p.
- ROMERO J., KAGALOU I., IMBERGER J., HELA D., KOTTI M., BARTZOKAS A., ALBANIS T., EVMIRIDES N., KARKABOUNAS S., PAPAGIANNIS J. & BITHAVA A. 2002. Seasonal water quality of shallow and eutrophic Lake Pamvotis, Greece: Implications for restoration. *Hydrobiologia*, 474: 91-105.
- ROUSKAS J. 2001. *The return of the Lake Karla*, unknown Publisher, Athens (in Greek), pp 186.
- SIDIROPOULOS P., MYLOPOULOS N. & LOUKAS A. 2015. Stochastic Simulation and Management of an Over-Exploited Aquifer Using an Integrated Modeling System. *Journal of Water Resources Management* 29(3): 929-943.
- ZALIDIS G, TAKAVAKOGLOU V., PANORAS A., BILAS G. & KATSAVOUNI S. 2004. Re-establishing a sustainable wetland at former Lake Karla, Greece, using Ramsar restoration guidelines. *Environmental Management* 34: 875-886.

Received: 22.11.2018
Accepted: 03.02.2019