

# Diet Composition of the Blotched Picarel *Spicara flexuosa* Rafinesque, 1810 (Actinopterygii: Sparidae) from the Aegean Sea, Turkey

*Tuncay Murat Sever*

Faculty of Fisheries, Ege University, 35100 Bornova-İzmir, Turkey; E-mail: tmsever66@gmail.com

**Abstract:** Stomach content of *Spicara flexuosa* from the Sığacık Bay (400 individuals) was studied. Overall, 64 taxa were identified belonging to 16 prey groups. The relative importance analysis identified crustacean groups (mostly pelagic copepods) as the most important prey in the diet of all size groups in all seasons. In addition, larvae of Decapoda and Mysidacea were also important in certain seasons and size groups. Based on the results of the study, *S. flexuosa* is a pelagic zooplanktivorous species that feeds with meso- and macrozooplankton, with pelagic copepods being the main food item for this species.

**Key words:** *Spicara flexuosa*, diet, zooplanktivore, Sığacık Bay.

## Introduction

Species of the genus *Spicara* Rafinesque, 1810 are schooling littoral fishes occurring in the Eastern Atlantic, throughout the Mediterranean Sea and part of the Black Sea. Although their commercial value is relatively low, they represent one of the major components of the Turkish coastal fisheries. Currently, there are indications of overfishing of *Spicara* spp. stocks in Turkey, which annual production has drastically decreased from a total of 2,250 tonnes in 2001 to 329 tonnes in 2016, with a mean production of 770 tonnes during the last decade (TUIK 2016).

The taxonomical discrimination between *Spicara maena* and *S. flexuosa* has long been a matter of dispute, primarily due to distinct colour changes reinforced by protogynous hermaphrodite sexual inversion (MINOS et al. 2013). Previous research on karyology (VASILIEV 1980), genetics (IMSIRIDOU et al. 2011, TURAN 2011, BEKTAS et al. 2018) and morphology (MINOS et al. 2013) have undoubtedly proved that *S. maena* and *S. flexuosa* are distinct species. This has also been accepted by renowned databases such as the Catalog of Fishes (ESCHMEYER et al. 2017) and World Register of Marine Species (WORMS 2017).

*Spicara flexuosa* Rafinesque, 1810 (blotched picarel) is distributed widely along the Turkish coastline (BILECENOĞLU et al. 2014). It prefers sandy or muddy bottoms in the coastal shelf up to 130 m (FISCHER et al. 1987). There are several studies on the biology of the species in the Mediterranean: age and growth – Ionian Sea (MYTILINEOU 1987, MYTILINEOU & PAPACONSTANTINO 1991), Aegean Sea (ARAMPATZI-ZIAMOU et al. 2011), Tunisian coast (HATTOUR et al. 1985), north-eastern Mediterranean Sea (ÖZVAROL 2014); genetics and morphology – Black Sea (VASILIEV 1980; VASILIEVA & SALEKHOVA 1983), Aegean Sea (İLKYAZ et al. 2007, IMSIRIDOU et al. 2011, TURAN 2011, MINOS et al. 2013, BEKTAŞ et al. 2018); reproduction – Ionian Sea (MYTILINEOU 1987). However, little work has been done on the feeding habits of the species, with examples available from the Ionian Sea (MYTILINEOU 1987) and the Black Sea (LIPSKAYA & SALEKHOVA 1980).

The aim of the present study was to examine the feeding habits of the blotched picarel from the Sığacık Bay of the central Aegean Sea by providing detailed information on the diet of the species based on different size groups and seasons and using qualitative and quantitative analyses.

## Materials and Methods

Blotched picarels were caught seasonally in the Sığacık Bay between May 2005 and June 2006 (Fig. 1) with the F/V Hapuloğlu commercial trawler. Sampling in the Sığacık Bay was at 100-200 m depth. After length measurements were done, the stomachs were removed and preserved in 4% formalin solution. At the laboratory, stomach content was examined and identified at the lowest possible taxonomic level group or at the species level using SZX7 Olympus stereomicroscope. Body parts that were not digested were used to count the prey organisms in stomachs (KATAĞAN 1990). Species identification for copepods was done following ROSE (1933), BRODSKII (1967) and RAZOULS et al. (2017). Items separated by either group or species were weighted as wet weight with precision of 0.0001 g sensitivity after being left on dry blotting paper for 10 min.

Quantitative representation of the stomach contents and estimation of the relative importance of different preys were done by using relative number of total gut content (%N), frequency of occurrence (%F) and relative weight (%W) (HYSLOP 1980). The participation of the main food items was estimated using the Index of Relative Importance (IRI);  $IRI = \%F * (\%N + \%W)$  (PINKAS et al. 1971). It was calculated and expressed as a percentage:  $IRI\% = 100 * IRI * (\sum IRI)^{-1}$  (CORTES 1998). The values were estimated for both seasonal and length groups, which were selected as  $\leq 11.9$  cm, 12.0-13.9 cm, 14.0-15.9 cm and  $\geq 16.0$  cm according to the length distribution in the samples.

The similarities between seasons and length groups were determined using the Bray–Curtis Similarity Index based on IRI% data and significant groups were identified using the SIMPROF test (WASHINGTON 1984, CLARKE et al. 2014).

## Results

Stomachs of 400 individuals of *S. flexuosa* (total length (TL) range=9.6–16.8 cm) were examined from the Sığacık Bay. Sixty-one stomachs were empty (15.25%): six from spring, 18 from summer, 16 from autumn and 21 from winter. According to the results of the analyses, 16 prey groups were identified belonging to eight taxonomic groups: Siphonophora, Polychaeta, Crustacea, Mollusca, Chaetognatha, Appendicularia, Thaliacea and Actinopterygii.

The pelagic copepods were the most important prey item in terms of IRI%, N%, F% and W% (in summer with Mysidacea for the 3rd and 4th



Fig. 1. Study area and stations.




Fig. 2. Dendrogram showing seasonal similarities in stomach contents of *Spicara flexuosa* based on the Bray-Curtis Index in the Sığacık Bay.




Fig. 3. Dendrogram showing size similarities in stomach contents of *Spicara flexuosa* based on the Bray-Curtis Index in the Sığacık Bay.

length groups) for all seasons and length groups in the Sığacık Bay (Tables 1 and 2). The remainder of the diet composition, on the other hand, varied between seasons and length groups. It consisted, in different ratios, of Cladocera, larvae of Decapoda, Bivalvia, Appendicularia, larvae of Brachyura, Chaetognatha, Cirripedia, Gastropoda, fish larvae and eggs, Siphonophora, Ostracoda, Stomatopoda, Thaliacea and Polychaeta. As a result, 64 taxa were found (Tables 1 and 2).

The abundance of pelagic copepods, which were the main prey in the diet of the blotched picarel (with 46 species) in the Sığacık Bay, exhib-

**Table 1.** Diet composition of *Spicara flexuosa* in the Sığacık Bay in sampling periods

	Spring (n=96)				Summer (n=77)				Autumn (n=66)				Winter (n=100)			
	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%
<b>Siphonophora</b>	0.20	0.69	25.00	0.20					0.58	2.81	33.33	0.81	0.50	1.69	10.00	0.24
<b>Polychaeta</b>					0.79	24.27	10.39	1.70	0.15	3.12	16.67	0.39				
<b>Crustacea</b>																
<b>Cladocera</b>	0.17	0.17	8.33	0.02									4.31	2.28	35.00	2.41
<i>Penilia avirostris</i>													0.59	0.64	12.00	0.16
<i>Podon intermedius</i>	0.17	0.17	8.33	0.02									3.72	1.64	30.00	1.77
<b>Ostracoda</b>	0.68	0.41	41.67	0.40					6.56	2.11	83.33	5.17				
<b>Copepoda</b>	81.12	58.45	100.00	83.03	19.43	4.80	37.66	5.82	87.63	80.98	98.48	91.65	82.11	48.04	65.00	88.26
Calanoida	13.13	7.76	100.00	18.25	2.06	0.61	28.57	0.50	8.01	3.52	98.48	8.13	5.63	3.33	65.00	6.40
<i>Acartia clausi</i>													1.45	1.58	21.00	0.70
<i>Acartia negligens</i>													7.49	5.73	33.00	4.80
<i>Nannocalanus minor</i>	0.88	1.20	25.00	0.46					2.34	3.12	50.00	1.96				
<i>Neocalanus gracilis</i>	0.07	0.13	16.67	0.03												
<i>Neocalanus tenuicornis</i>									0.16	0.82	16.67	0.12				
<i>Paracalanus parvus</i>	3.26	2.39	58.33	2.88	0.88	0.32	11.69	0.09					0.54	0.53	11.00	0.13
<i>Paracalanus nanus</i>	0.10	0.21	8.33	0.02												
<i>Clausocalanus arcuicornis</i>	4.08	1.75	66.67	3.40	0.98	0.39	11.69	0.10	1.02	1.38	66.67	1.15				
<i>Clausocalanus furcatus</i>	3.88	2.29	58.33	3.14					0.72	1.63	50.00	0.84	0.50	0.64	12.00	0.15
<i>Clausocalanus</i> spp.	19.29	14.25	66.67	19.53					0.73	0.98	68.18	0.84	1.45	1.99	37.00	1.40
<i>Mecynocera clausi</i>	0.07	0.12	16.67	0.03					1.05	1.80	66.67	1.36				
<i>Calocalanus pavo</i>	0.03	0.05	8.33	0.01												
<i>Ctenocalanus vanus</i>	0.10	0.20	16.67	0.04												
<i>Ischnocalanus plumulosus</i>	0.10	0.19	16.67	0.04					0.30	0.68	16.67	0.12				
<i>Temora stylifera</i>	11.94	6.76	41.67	6.81					0.16	0.28	18.18	0.06	0.95	0.94	20.00	0.42
<i>Pleuromamma abdominalis</i>	0.41	0.78	33.33	0.35					4.00	3.74	66.67	3.70				
<i>Pleuromamma gracilis</i>	0.44	0.87	33.33	0.38					2.66	6.33	83.33	5.37				
<i>Candacia armata</i>	0.65	1.70	50.00	1.02					0.17	1.46	18.18	0.21				
<i>Candacia simplex</i>	0.92	2.26	41.67	1.16					0.44	2.76	33.33	0.76				
<i>Centropages typicus</i>	4.59	2.26	50.00	2.99									59.58	30.04	71.00	69.95
<i>Lucicutia flavicornis</i>	1.29	1.02	50.00	1.01	0.98	0.41	9.09	0.08	13.88	7.26	81.82	12.39				
<i>Lucicutia</i> spp.									0.33	1.29	16.67	0.19				
<i>Haloptilus longicornis</i>	0.10	0.30	8.33	0.03					4.66	2.17	43.94	2.15				
<i>Scolecithrix bradyi</i>									0.15	0.39	15.15	0.06				
<i>Diaixis pygmoea</i>	0.03	0.08	8.33	0.01												
<i>Euaetideus giesbrechti</i>									0.32	0.68	15.15	0.11				
<i>Isias clavipes</i>	0.14	0.26	16.67	0.06	0.88	0.17	10.39	0.07								
<i>Corycaeus giesbrechti</i>	0.58	0.57	50.00	0.50					0.19	0.14	18.18	0.04				
<i>Corycaeus typicus</i>	2.31	1.27	66.67	2.09	6.97	1.41	37.66	2.05	7.74	5.40	100.00	9.41	0.50	0.64	12.00	0.15
<i>Corycaeus flaccus</i>	0.20	0.24	16.67	0.06					2.50	1.91	66.67	2.11				
<i>Ditrichocorycaeus brehmi</i>	0.65	0.42	33.33	0.31					0.73	1.18	50.00	0.68	0.95	0.70	20.00	0.36
<i>Corycaeus clausi</i>	0.20	0.25	16.67	0.07					1.91	2.14	50.00	1.45				
<i>Corycaeus ovalis</i>	0.41	0.26	41.67	0.24												
<i>Corycaeus lautus</i>	0.07	0.10	16.67	0.02												
<i>Urocorycaeus furcifer</i>	0.44	0.49	33.33	0.27					2.49	3.12	50.00	2.01				
<i>Farranula rostrata</i>	0.71	0.35	58.33	0.54	4.12	0.80	19.48	0.62	0.16	0.20	16.67	0.04	0.95	0.58	20.00	0.34

Table 1. Continuation.

	Spring (n=96)				Summer (n=77)				Autumn (n=66)				Winter (n=100)			
	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%
<i>Corycaeus</i> spp.	0.17	0.19	33.33	0.11					0.29	0.98	33.33	0.30				
<i>Vetтория parva</i>									0.45	0.51	33.33	0.23				
<i>Oncaea media</i>	5.51	2.49	75.00	5.24	0.88	0.13	11.69	0.08	3.79	1.21	83.33	2.98	1.50	0.64	31.00	0.73
<i>Oncaea mediterranea</i>	2.18	1.75	58.33	2.00	0.88	0.37	10.39	0.08	5.99	2.25	83.33	4.92				
<i>Oncaea venusta</i>	0.24	0.44	33.33	0.20												
<i>Triconia conifera</i>	0.85	0.66	50.00	0.66	0.79	0.19	9.09	0.06	10.97	7.26	100.00	13.05				
<i>Oithona plumifera</i>	0.07	0.18	16.67	0.04					0.87	1.60	66.67	1.18	0.59	0.70	12.00	0.17
<i>Lubbockia squillimana</i>	0.34	0.62	25.00	0.21					0.60	1.24	36.36	0.48				
<i>Lubbockia aculeata</i>	0.07	0.18	16.67	0.04					1.30	2.84	51.52	1.53				
<i>Pontoeciella abyssicola</i>									0.15	0.62	16.67	0.09				
<i>Sapphirina metallina</i>	0.54	1.09	58.33	0.83					5.67	6.84	98.48	8.82				
<i>Microsetella rosea</i>	0.07	0.08	8.33	0.01					0.44	0.73	30.30	0.25				
<i>Microsetella gracilis</i>									0.16	0.25	15.15	0.04				
Harpacticoida									0.15	0.28	16.67	0.05				
<i>Lucifer</i> sp.	0.24	0.31	8.33	0.04												
<b>Cirripedia nauplii</b>													0.50	0.23	11.00	0.09
<b>Cypris larvae</b>	0.07	0.04	16.67	0.02					0.15	0.20	16.67	0.04	1.36	0.82	11.00	0.26
<b>Mysidacea</b>	5.48	32.17	58.33	19.19	79.00	69.82	97.40	94.45								
<b>Stomatopoda</b>									0.16	2.87	18.18	0.39				
<b>Decapoda larvae</b>	1.02	0.81	83.33	1.33									1.45	23.85	21.00	5.84
<b>Brachyura larvae</b>	0.92	1.90	41.67	1.03									0.50	15.08	10.00	1.71
<b>Mollusca</b>																
<b>Gastropoda</b>	8.23	1.84	16.67	1.47					1.74	2.11	16.67	0.46				
<b>Bivalvia</b>	0.03	0.04	8.33	0.01												
<b>Chaetognatha</b>	0.78	0.76	33.33	0.45									5.09	4.50	11.00	1.16
<b>Appendicularia</b>	0.92	0.33	50.00	0.55					1.75	0.79	68.18	1.24	3.72	2.81	41.00	2.94
<b>Thaliacea</b>																
<i>Salpa</i> sp.					0.79	1.10	9.09	0.11	1.30	5.01	51.52	2.33				
<b>Teleostei</b>																
Fish larvae	0.03	1.70	8.33	0.13												
Fish eggs	0.10	0.35	25.00	0.10									0.45	0.70	10.00	0.13

ited seasonal changes. The dominant food items were: in spring – *Temora stylifera*, *Oncaea media*, *Centropages typicus*, *Clausocalanus arcuicornis* and *Clausocalanus furcatus*; in summer – *Corycaeus typicus* and *Farranula rosata*; in autumn – *Lucicutia flavicornis*, *Triconia conifera*, *Corycaeus typicus*, *Oncaea mediterranea*, *Sapphirina metallina*, *Haloptilus longicornis* and *Pleuromamma abdominalis*; in winter – *Centropages typicus*, *Acartia negligens* and *O. media* (Table 1).

According to the Bray-Curtis cluster analysis, the highest similarity was detected between summer and winter (40.56%; Fig. 2) and between the 1st and 2nd length groups (58.71%; Fig. 3). These groupings were statistically significant (SIMPROF test,  $P < 0.05$ ).

## Discussion

In the Mediterranean, stomach content studies on blotched picarel' are limited. LIPSKAYA & SALEKHOVA (1980) have reported that, even though copepods are dominant in the Black Sea, in blotched picarel stomach contents mysids, amphipods and decapods have also been observed in high ratios. MYTILINEOU (1987) has reported that the species in the Gulf of Patraikos is feeding on zooplankton and copepods are dominant. He has also indicated that *S. flexuosa* feeds more between September and November than during the other months.

Two studies on this species from the Mediterranean, by STERGIOU & KARPOUZI (2002) on 332 species and by NAVARRO et al. (2009) near

**Table 2.** Diet composition of *Spicara flexuosa* in the Sığacık Bay by length groups.

	10.0-11.9 cm (n=72)				12.0-13.9 cm (n=126)				14.0-15.9 cm (n=91)				≥16.0 cm (n=50)			
	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%
<b>Siphonophora</b>	0.57	2.50	16.67	0.63	0.14	0.59	19.05	0.18	8.89	1.58	14.29	1.33				
<b>Polychaeta</b>									0.29	2.34	14.29	0.33	0.79	58.53	20.00	18.01
<b>Crustacea</b>																
<b>Cladocera</b>	0.42	0.38	15.28	0.09	0.39	0.35	14.29	0.08								
<i>Penilia avirostris</i>					0.04	0.07	4.76	0.01								
<i>Podon intermedius</i>	0.42	0.38	15.28	0.15	0.35	0.28	14.29	0.12								
<b>Ostracoda</b>	0.14	0.10	15.28	0.05	1.77	0.74	33.33	1.10	3.82	0.46	14.29	0.54	0.87	0.05	20.00	0.28
<b>Copepoda</b>	96.61	87.99	70.83	95.57	79.48	60.74	80.95	87.23	47.61	17.75	57.14	27.68	67.33	19.72	40.00	45.45
Calanoida	15.83	12.33	65.28	22.55	11.24	6.49	80.95	18.77	4.41	0.91	57.14	2.70	9.57	1.56	40.00	6.76
<i>Acartia clausi</i>					0.11	0.18	9.52	0.04								
<i>Acartia negligens</i>	0.57	0.87	31.94	0.56	0.43	0.48	4.76	0.06								
<i>Nannocalanus minor</i>	0.42	0.76	15.28	0.22	1.03	1.36	19.05	0.59	2.94	1.45	14.29	0.56				
<i>Neocalanus gracilis</i>	0.14	0.35	15.28	0.09	0.04	0.07	4.76	0.01								
<i>Neocalanus tenuicornis</i>					0.04	0.19	4.76	0.01								
<i>Paracalanus parvus</i>	3.84	3.72	48.61	4.51	2.52	1.89	28.57	1.65								
<i>Paracalanus nanus</i>					0.11	0.21	4.76	0.02								
<i>Clausocalanus arcuicornis</i>	0.85	0.49	33.33	0.55	4.25	2.08	42.86	3.55	0.59	0.15	16.48	0.11	0.87	0.10	20.00	0.30
<i>Clausocalanus furcatus</i>	1.43	1.04	15.28	0.46	3.79	2.42	38.10	3.09	0.59	0.46	15.38	0.14	0.79	0.21	20.00	0.30
<i>Clausocalanus</i> spp.	0.85	1.35	65.28	1.76	20.10	14.67	42.86	19.48					1.74	0.57	42.00	1.48
<i>Mecynocera clausi</i>	0.14	0.31	15.28	0.09	0.21	0.35	19.05	0.14	0.59	0.40	14.29	0.13				
<i>Calocalanus pavo</i>					0.04	0.05	4.76	0.01								
<i>Ctenocalanus vanus</i>					0.11	0.20	9.52	0.04								
<i>Ischnocalanus plumulosus</i>	0.14	0.31	13.89	0.08	0.14	0.29	9.52	0.05								
<i>Temora stylifera</i>	44.53	32.78	31.94	30.29	1.38	0.88	28.57	0.84								
<i>Pleuromamma abdominalis</i>	0.28	0.69	16.67	0.20	0.96	1.17	28.57	0.80	3.53	1.16	13.19	0.55				
<i>Pleuromamma gracilis</i>	0.57	1.42	15.28	0.37	0.74	1.55	28.57	0.86	1.47	1.24	14.29	0.34	1.66	1.25	24.00	1.06
<i>Candacia armata</i>	1.27	4.34	13.89	0.96	0.39	1.25	23.81	0.51								
<i>Candacia simplex</i>	0.71	2.26	30.56	1.11	0.82	2.09	19.05	0.72	0.59	1.39	14.29	0.25				
<i>Centropages typicus</i>	8.60	7.78	63.89	12.84	7.13	4.20	42.86	6.35								
<i>Lucicutia flavicornis</i>	0.99	1.01	15.28	0.37	3.08	1.95	42.86	2.82	7.94	1.54	14.29	1.20	10.36	1.88	22.00	4.09
<i>Lucicutia</i> spp.									0.59	0.97	13.19	0.18				
<i>Haloptilus longicornis</i>					0.92	0.66	14.29	0.30	2.65	0.46	15.38	0.43				
<i>Scolecithrix bradyi</i>													0.87	0.73	18.00	0.44
<i>Diaixis pygmoea</i>					0.04	0.08	4.76	0.01								
<i>Euaetideus giesbrechti</i>													1.66	1.25	18.00	0.80
<i>Isias clavipes</i>	0.28	0.69	13.89	0.17	0.07	0.14	4.76	0.01	0.29	0.17	14.29	0.06				
<i>Corycaeus giesbrechti</i>	0.28	0.35	29.17	0.23	0.57	0.54	23.81	0.35								
<i>Corycaeus typicus</i>	2.54	1.81	27.78	1.48	3.30	2.21	61.90	4.46	3.82	1.01	42.86	1.84	6.49	1.51	24.00	2.91
<i>Corycaeus flaccus</i>					0.46	0.43	19.05	0.22	0.88	0.25	14.29	0.14	5.62	1.46	18.00	1.93
<i>Ditrichocorycaeus brehmi</i>	1.70	1.53	33.33	1.32	0.50	0.49	33.33	0.43								
<i>Corycaeus clausi</i>					0.46	0.52	14.29	0.18	0.59	0.25	14.29	0.11	3.24	1.20	20.00	1.35
<i>Corycaeus ovalis</i>	0.57	0.45	18.06	0.23	0.28	0.18	19.05	0.12								
<i>Corycaeus lautus</i>					0.07	0.10	9.52	0.02								
<i>Urocorycaeus furcifer</i>	0.42	0.63	16.67	0.21	0.89	1.03	23.81	0.60	0.59	0.27	14.29	0.11				
<i>Farranula rostrata</i>	1.55	1.18	62.50	2.10	0.53	0.33	28.57	0.32	0.88	0.38	14.29	0.16	0.95	0.37	22.00	0.44

Table 2. Continuation.

	10.0-11.9 cm (n=72)				12.0-13.9 cm (n=126)				14.0-15.9 cm (n=91)				≥16.0 cm (n=50)			
	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%	N%	W%	F%	IRI%
<i>Corycaeus</i> spp.	0.14	0.21	15.28	0.07	0.18	0.27	19.05	0.11					0.87	0.94	24.00	0.66
<i>Vetoria parva</i>					0.11	0.12	9.52	0.03								
<i>Oncaea media</i>	3.86	3.78	70.83	6.65	4.96	2.10	42.86	3.96	0.88	0.19	42.86	0.41	4.11	0.47	24.00	1.67
<i>Oncaea mediterranea</i>	2.69	2.81	34.72	2.34	2.59	1.72	42.86	2.41	2.35	0.34	14.29	0.34	4.83	0.57	22.00	1.80
<i>Oncaea venusta</i>	0.42	1.01	13.89	0.24	0.14	0.26	14.29	0.08								
<i>Triconia confifera</i>	0.57	0.56	16.67	0.23	2.52	1.68	38.10	2.09	6.76	1.77	27.47	2.08	2.37	0.63	40.00	1.82
<i>Oithona plumifera</i>	0.14	0.42	18.06	0.12	0.21	0.44	19.05	0.16	0.29	0.19	15.38	0.07	0.87	0.47	22.00	0.45
<i>Lubbockia squillimana</i>					0.46	0.85	19.05	0.33	0.29	0.23	14.29	0.07				
<i>Lubbockia aculeata</i>					0.25	0.55	19.05	0.20	1.18	0.95	14.29	0.27				
<i>Pontoeciella abyssicola</i>					0.04	0.14	4.76	0.01								
<i>Sapphirina metallina</i>	0.28	0.76	13.89	0.18	1.21	1.72	38.10	1.46	2.06	1.03	28.57	0.79	9.57	4.12	38.00	7.90
<i>Microsetella rosea</i>					0.07	0.09	4.76	0.01	0.59	0.38	13.19	0.11	0.87	0.42	20.00	0.39
<i>Microsetella gracilis</i>									0.29	0.19	14.29	0.06				
Harpacticoida					0.04	0.07	4.76	0.01								
<i>Lucifer</i> sp.													5.62	2.50	20.00	2.47
Cirripedia nauplii					0.04	0.03	4.76	0.01								
Cypris larvae					0.18	0.14	14.29	0.05	0.29	0.15	13.19	0.05				
<b>Mysidacea</b>	0.14	1.08	13.89	0.21	4.64	25.73	33.33	13.24	34.98	74.81	85.71	83.68	19.78	3.81		35.81
<b>Stomatopoda</b>					0.04	0.67	4.76	0.04								
<b>Decapoda larvae</b>	1.13	5.69	63.89	5.35	0.74	2.32	28.57	1.15	0.29	0.08	13.19	0.04	2.45	0.63	20.00	0.93
<b>Brachyura larvae</b>	0.57	1.53	31.94	0.82	0.82	3.27	14.29	0.76					0.79	0.57	20.00	0.41
<b>Mollusca</b>																
<b>Gastropoda</b>					9.00	2.37	19.05	2.83								
<b>Bivalvia</b>					0.04	0.05	4.76	0.01								
<b>Chaetognatha</b>					1.17	1.25	19.05	0.60					0.79	0.26	24.00	0.38
<b>Appendicularia</b>	0.28	0.10	16.67	0.08	1.31	0.70	52.38	1.38	2.06	0.34	14.29	0.30	0.79	0.10	20.00	0.27
<b>Thaliacea</b>																
<i>Salpa</i> sp.					0.14	0.73	14.29	0.16	1.76	2.49	13.19	0.50				
<b>Teleostei</b>																
Fish larvae													0.79	13.82	22.00	4.88
Fish eggs	0.14	0.63	16.67	0.16	0.11	0.32	14.29	0.08								

the Balearic Islands on 16 species reported results on feeding preferences of the blotched picarel. Both studies have shown that this fish species is planktivorous.

The present study shows that the food composition of *S. flexuosa* is relatively rich and includes diverse zooplanktonic groups and species. Nonetheless, certain groups are preferred: pelagic copepods (for all seasons) and mysids (for some seasons). Feeding habits of planktivorous fishes are affected directly by their behaviour and food quantity in their habitat (MAURIN 1966). HELA & LAEVAUSTU (1961, cited after FURNESTIN et al. 1966) have reported that pelagic fish species are abundant in a zone where phytoplankton and zooplankton are also abundant.

The size groups I and II of the blotched picarel feed on small-sized prey (copepods such as *Oncaea*

spp., *Corycaeus* spp. and *T. stylifera*, gastropods and small mysids), while the size groups III and IV prefer larger-sized prey (big mysids, larger copepods such as *Lucicutia flavicornis*, *Pleuromamma* spp. and others). As a conclusion, copepods were the most important prey group for all size groups of the species. For a planktivore fish, copepods are an indispensable source (SORBE 1972). The blotched picarel, being a benthopelagic fish, is known to generally feed on zooplankton species and occasionally on benthic invertebrates (COLL et al. 2006, 2008), which are less important source in trophic web (STERGIOU & KARPOUZI 2002, NAVARRO et al. 2009). Findings of our study show that the blotched picarel feeds on mostly holoplanktonic (especially copepods and mysids) and, to a smaller extent, on meroplanktonic

larvae (decapods, brachyurans and polychaetes).

A wide spectrum of prey items have been found in the stomachs of the blotched picarel in the Sığacık Bay. There are also rarely consumed prey items such as Isopoda, Stomatopoda, Bivalvia, Gastropoda, Thaliacea, Siphonophora and Chaetognatha.

Pelagic copepods have been reported as the dominant group by zooplankton studies from various areas of the Aegean Sea (PAVLOVA 1966, PASTEUR 1976, MATSAKIS et al. 1978, MORAITOU-APOSTOLOPOULOU 1985, SEVER 1997, 2009, BENLI et al. 2001, AKER 2002, İŞİNIBİLİR 2009). SEVER (1997) reported for the Aegean Sea that while *C. arcuicornis*, *C. typicus*, *C. giesbrechti*, *H. longicornis*, *F. rostrata*, *O. media* live in neritic and oceanic zone, *A. clausi* and *T. stylifera* are abundant only in the neritic zone. *C. typicus*, *O. media*, *A. clausi* and *T. stylifera* are dominant in the same area in all seasons (AKER 2002) but other species are seasonally abundant. In the Sığacık Bay, the abundant copepod species are *Centropages typicus*, *T. stylifera*, *Corycaeus typicus*, *L. flavicornis*, *T. conifera*, *A. negligens*, *H. longicornis*, *F. rostrata*, *O. media* and *C. arcuicornis*. The species found in stomachs of the examined fish individuals have been reported to be abundant in the zooplankton in the area by SEVER (1997) and AKER (2002).

According to the stomach content analyses, blotched picarel is identified as a meso-macro zooplankton feeder and pelagic zooplanktivore, with pelagic copepods being the most important prey group. In addition, this species is important food for benthic-pelagic predators such as *Merluccius merluccius*, *Synodus saurus* and *Zeus faber*, which occupy higher positions in the food web (CARPENTIERI et al. 2005, ESPOSITO et al. 2009; STERGIU & FOURTOUNI 1991). In this sense, it plays a very important role in the energy flow between the lower and upper trophic levels of benthic and pelagic food webs.

**Acknowledgement:** This study was supported by the Ege University Scientific Research Fund (Project No: 2005/SÜF/007).

## References

- AKER V. 2002. Seasonal Distribution of Planktonic Copepods in the Turkish Coastal Waters of the Middle Aegean Sea. PhD Thesis, Ege University, 107 p. [in Turkish].
- ARAMPATZI-ZIAMOU D., KARELLOU E.I.E., STAMAS A., CHASAN E., GOLOMAZOU E., NEOFITOU N., AUFANTI S. & NEOFITOU C. 2011. Study of Biological Characteristics of Picarel (*Spicara flexuosa* Rafinesque, 1810) in Pagasitikos Gulf. In: Fourth International Symposium "Hydrobiology – Fisheries", Volos, Greece, 9-11 June 2011, pp. 223-226. [In Greek].
- BEKTAŞ Y., AKSU İ., KALAYCI G., IRMAK E., ENGIN S. & TURAN D. 2018. Genetic differentiation of three *Spicara* (Pisces: Centranchidae) species, *S. maena*, *S. flexuosa* and *S. smaris*: an intraspecific substructure of *S. flexuosa* in Turkish coastal waters. Turkish Journal of Fisheries and Aquatic Sciences 18: 301-311. DOI 10.4194/1303-2712-v18\_2\_09.
- BENLI H.A., TARKAN A.N. & SEVER T.M. 2001. Comparison of the mesozooplankton composition the southwestern Black Sea, Sea of Marmara and Eastern Aegean Sea. Turkish Journal of Marine Sciences 7: 163-179.
- BILECENOĞLU M., KAYA M., CİHANGİR B. & ÇİÇEK E. 2014. An updated checklist of the marine fishes of Turkey. Turkish Journal of Zoology 38: 901-929.
- BRODSKII K.A. 1967. Calanoida of the Far-Eastern Seas and polar basin of the USSR. Keys to the Fauna of the USSR, 35. Israel Program for Scientific Translations, Jérusalem, Israel. 440 p.
- CARPENTIERI P., COLLOCA F., CARDINALE M., BELLUSCIO A. & ARDIZZONE G.D. 2005. Feeding habits of European hake (*Merluccius merluccius*) in the central Mediterranean Sea. Fishery Bulletin 103 (2): 411-416.
- CLARKE K.R., GORLEY R.N. & WARWICK R.M. 2014. Change in marine communities: an approach to statistical analysis and interpretation, third edition. PRIMER-E Ltd., Plymouth, UK.
- COLL M., PALOMERA I. & TUDELA S. 2006. Trophic flows, ecosystem structure and fishing impacts in the South Catalan Sea, North western Mediterranean. Journal of Marine Systems 59: 63–96.
- COLL M., PALOMERA I. & TUDELA S. 2008. Food-web dynamics in the South Catalan Sea ecosystem (NW Mediterranean) for 1978–2003. Ecological Modelling 217: 95–116. DOI: 10.1016/j.ecolmodel.2008.06.013.
- CORTÉS E. 1998. Methods of studying fish feeding: reply. Canadian Journal of Fisheries and Aquatic Sciences 55 (12): 2709. <https://doi.org/10.1139/f98-202>.
- COSTA F. 1991. Atlante dei pesci dei mari italiani. Gruppo Ugo Mursia Editore S.p.A. Milano, Italy. 438 p.
- ESCHMEYER W.N., FRICKE R. & VAN DER LAAN R. (Eds.) 2017. Catalog of fishes: Genera, Species, References. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. Electronic version accessed December 2017.
- ESPOSITO V., BATTAGLIA P., CASTRIOTA L., FINOIA M.G., SCOTTI G. & ANDALORO F. 2009. Diet of Atlantic lizardfish, *Synodus saurus* (Linnaeus, 1758) (Pisces: Synodontidae) in the central Mediterranean Sea. Scientia Marina 73 (2): 369-376. DOI: 10.3989/scimar.2009.73n2369.
- FISCHER W., BAUCHOT M.-L. & SCHNEIDER M. 1987. Fiches FAO d'identification des espèces pour les besoins de la pêche. (Révision 1). Méditerranée et mer Noire. Zone de pêche 37. Vol. II. Vertébrés, FAO, Rome, pp. 1031-1036.
- FROESE R. & PAULY D. 2017. FishBase version (08/2017). World Wide Web electronic publication. <http://www.fishbase.org> (20/08/2017).
- FURNESTIN M.L., MAURIN C., LEE J.Y. & RAIMBAULT R. 1966. Éléments de Planctologie Appliquée. Revue des Travaux de l'Institut des Pêches Maritimes, Paris 30(2): 117-278.
- GOLANI D., ÖZTÜRK B. & BAŞUSTA N. 2006. Centranchidae. In: Fishes of the Eastern Mediterranean, pp. 168-169. Turkish Marine Research Foundation, Istanbul.
- HATTOUR A., TURKI B. & ZAMMOURI N. 1985. Quelques aspects de la biologie de l'espèce *Spicara flexuosa* (Rafinesque, 1810) des eaux Tunisiennes. Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche de Salammbô 12: 143-162.
- HELA I. & LAEVASTU T. 1961. Fisheries hydrography. Fishing News (Books), London, 137 p.

- HYSLOP E.J. 1980. Stomach content analysis – a review of methods and their applications. *Journal of Fish Biology* 17: 411-429. DOI: 10.1111/j.1095-8649.1980.tb02775.x.
- İŞİNİBİLİR M. 2009. Summer mesozooplankton communities in the Turkish coastal waters of north Aegean Sea. *Journal of Fisheries Sciences.com* 3 (3): 237-242. DOI: 10.3153/jfsc.com.2009.029.
- ILKYZAZ A.T., METİN G. & KINACIGİL H.T. 2007. Using morphological differences of *Spicara* genus (*Spicara smaris*, *S. flexuosa* and *S. maena*) for species identification in the Aegean Sea. Fourteenth National Fisheries Symposium, 4-7 September 2007, Muğla. [In Turkish].
- İMSİRİDOU A., MINOS G., GAKOPOULOU A., KATSARES V., KARIDAS T.H. & KATSELIS G. 2011. Discrimination of two picarel species *Spicara flexuosa* and *Spicara maena* (Pisces: Centranchthidae) based on mitochondrial DNA sequences. *Journal of Fish Biology* 78: 373-377.
- KATAĞAN T., KAYA M., ERGEN Z. & ÖNEN M. 1990. İzmir Körfezinde Yaşayan *Mullus barbatus* Linnaeus, 1758 Türünün Beslenme Rejimi Doğa-Tr. [Feeding regime of *Mullus barbatus* Linnaeus, 1758 in İzmir Bay. Doğa-Tr.] *Journal of Zoology* 14 (2): 179-187.
- LIPSKAYA N. Y. & SALEKHOVA L. P. 1980. Studies in nutrition and Morphophysiological indicators of the genus *Spicara* fishes. *Ekologiya Morya* 2: 66-81. [In Russian].
- MAURIN C. 1966. Les poissons planctonophages. I. Caracteres Generaux, Comportement. In: *Éléments de planctologie appliqué*. Revue des Travaux de l'Institut des Pêches Maritimes, Paris 30 (2): 209-216.
- MINOS G., İMSİRİDOU A. & KATSELIS G. 2013. Use of morphological differences for the identification of two picarel species *S. flexuosa* and *S. maena* (Pisces: Centranchthidae). *Mediterranean Marine Science* 14 (3): 26-31. <http://dx.doi.org/10.12681/mms.423>.
- MYTILINEOU C. 1987. Contribution to the Biology of picarel *Spicara flexuosa* (Raf. 1810) in the Patraikos Gulf (Greece) (in Greek). Ph.D. Thesis, University of Athens, 152 p.
- MYTILINEOU C. & PAPAConstantinou C. 1991. Age and growth of *Spicara flexuosa* (Rafinesque, 1810) (Pisces, Centranchthidae) in the Patraikos Gulf (Greece). *Scientia Marina* 55 (3): 483-490.
- MATSAKIS J., YANNOPOULOS C., THESSALOU M., HATZAKIS A., MALOUHOU-GRIMBA V., MORAITOU-APOSTOLOPOULOU M., MYLONAS M., SIOKOU J. & YANNOPOULOS A. 1978. Ecologie marine de la région de l'Eubée du nord. III. Évolution du zooplancton dans sept stations. *Biologia Gallo-Hellenica* 7 (1-2): 99-144.
- MORAITOU-APOSTOLOPOULOU M. 1985. The zooplankton communities of the Eastern Mediterranean (Levantine Basin, Aegean Sea); influence of man made factors. In: MORAITOU-APOSTOLOPOULOU M. & KIORTSIS V. (Eds.) *Mediterranean Marine Ecosystems*. NATO Conference Series (I Ecology), Vol. 8. Springer, Boston, MA, pp. 303-329. DOI: 10.1007/978-1-4899-2248-9.
- NAVARRO J., LOUZAO M. & IGUAL J. M. 2009. Seasonal changes in the diet of a critically endangered seabird and the importance of trawling discards. *Marine Biology* 156: 2571–2578. DOI: 10.1007/s00227-009-1281-3.
- ÖZVAROL Y. 2014. Length-weight relationships of 14 fish species from the Gulf of Antalya (north-eastern Mediterranean Sea, Turkey). *Turkish Journal of Zoology* 38: 342-346. DOI: 10.3906/zoo-1308-44.
- PAPAConstantinou C. 1988. Check-list of marine fishes of Greece. National Centre for Marine Research, Hellenic Zoological Society, Athens. 257 p.
- PAVLOVA E. 1966. Composition and distribution of zooplankton in the Aegean Sea. In: *Investigation of plankton in South Seas* 7: 38-61.
- PASTEUR R., BERDUGO V. & KIMOR B. 1976. The abundance, composition and seasonal distribution of epizooplankton in coastal and offshore waters of the Eastern Mediterranean. *Acta Adriatica* 18 (4): 53-80.
- PINKAS L., OLIPHANT M. S. & IVERSON I. L. K. 1971. Food habits of albacore, bluefin tuna and bonito in California waters. State of California The Resources Agency, Department of Fish and Game, Fish Bulletin 152: 1–105.
- RAZOULS C., DE BOVÉE F., KOUWENBERG J. & DESREUMAUX N. 2017. Diversity and geographic distribution of marine planktonic copepods. <http://copepodes.obs-banyuls.fr/en> (05/10/2017).
- ROSE M. 1933. Copépodes pélagiques. Faune de France. Vol. 26. Fédération Française des sociétés de sciences naturelles; office central de faunistique. Paul Lechevalier, Paris.
- SEVER T.M. 1997. Establishment of pelagic copepods and quantitative and qualitative distributions of important copepod species in Aegean sea of Turkey. PhD Thesis, University of Dokuz Eylül, 133 p. [In Turkish].
- SEVER T.M. 2009. Pelagic Copepoda fauna of the Aegean Sea and the distribution of the common species. E.Ü. *Journal of Fisheries & Aquatic Sciences* 26 (3): 203-209.
- SORBE J.C. 1972. Écologie et éthologie alimentaire de l'ichthyofaune chlutable du plateau continental Sud Gascogne. Thèse de Doct. 3ème cycle, Aix-Marseille II, 125 p.
- STERGIOU K.I. & FOURTOUNI H. 1991. Food habits, ontogenetic diet shift and selectivity in *Zeus faber* Linnaeus, 1758. *Journal of Fish Biology* 39 (4): 589-603. DOI: 10.1111/j.1095-8649.1991.tb04389.x.
- STERGIOU K.I. & KARPOUZİ V. 2002. Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries* 11: 217–254.
- TORTONESE E. 1986. Centranchthidae. In: WHITEHEAD P.J.P., BAUCHOT M.-L., HUREAU J.-C., NIELSEN J. & TORTONESE E. (Eds.): *Fishes of the north-eastern Atlantic and the Mediterranean*, Paris, France: UNESCO, pp. 908-911.
- TUİK 2016. Su Ürünleri İstatistikleri Veri Tabanı. [www.tuik.gov.tr/PreIstatistikTablo.doistab\\_id=693](http://www.tuik.gov.tr/PreIstatistikTablo.doistab_id=693)
- TURAN C. 2011. The systematic status of the Mediterranean *Spicara* species (Centranchthidae) inferred from mitochondrial 16s rDNA sequence and morphological data. *Journal of Black Sea/Mediterranean Environment* 17 (1): 14-31.
- VASILIEV V.P. 1980. The selective mortality of different chromosome morphs in *Spicara flexuosa* Raf. *Genetica* 52/53: 327-331.
- VASILIEVA E.D. & SALEKHOVA L.P. 1983. The precision of diagnoses of the *Spicara* (Perciformes, Centranchthidae) species, based on osteological features. *Zoological Journal* 62(7), 1044-1056. [In Russian].
- WASHINGTON H.G. 1984. Diversity, biotic and similarity indices, a review with special reference to aquatic ecosystems. *Water Research* 18: 653-694. DOI: 10.1016/0043-1354(84)90164-7.
- WORMS EDITORIAL BOARD 2017. World register of marine species. Available from <http://www.marinespecies.org> at VLIZ. Accessed 2017-12-20. DOI: 10.14284/170.

Received: 15.05.2018

Accepted: 12.01.2019